

Kinetic theory of fermions in curved spacetime

Christian Fidler^{1,*} and Cyril Pitrou^{2,†}

¹*Catholic University of Louvain - Center for Cosmology, Particle Physics and Phenomenology (CP3) 2,
Chemin du Cyclotron, B-1348 Louvain-la-Neuve, Belgium*

²*Institut d'Astrophysique de Paris, CNRS-UMR 7095, UPMC - Paris VI,
Sorbonne Universités, 98 bis Bd Arago, 75014 Paris, France*

(Dated: January 13, 2017)

We build a statistical description of fermions, taking into account the spin degree of freedom in addition to the momentum of particles, and we detail its use in the context of the kinetic theory of gases of fermions particles. We show that the one-particle distribution function needed to write a Liouville equation is a spinor valued operator. The degrees of freedom of this function are covariantly described by an intensity function and by a polarisation vector which are parallel transported by free streaming. Collisions are described on the microscopic level and lead to a Boltzmann equation for this operator. We apply our formalism to the case of weak interactions, which at low energies can be considered as a contact interaction between fermions only. This allows to discuss the structure of the collision term for a few typical weak-interaction mediated reactions. In particular we find for massive particles that a dipolar distribution of velocities in the interacting species is necessary to generate linear polarisation, as opposed to the case of photons for which linear polarisation is generated from the quadrupolar distribution of velocities.

Introduction

Out of equilibrium evolution of gases is well described in the context of kinetic theory. This description is based on a distribution function $f(t, \mathbf{x}, \mathbf{p})$, which describes the distribution of particles in phase space, that is at the position \mathbf{x} for the spatial momentum \mathbf{p} at a given time t . This description can be generalised to the relativistic context and is used in curved space-time when treating simultaneously gravitational effects [1, 2]. If collisions can be ignored and gas particles are free streaming, then the evolution of the distribution functions is dictated by a Liouville equation. However, when interactions, between the same species or among species, cannot be ignored, a collision term must be added and this leads to the Boltzmann equation dictating the evolution of the distribution function.

In many circumstances, this description needs to be approximated to reduce the computational cost. If interactions are very strong, the gas is driven toward local thermal equilibrium and behaves effectively as a fluid with pressure. Thus, when decomposing the directional dependence in the momentum, one needs only the monopole which is related to the energy density of the fluid, and the dipole which is related to the fluid velocity. If interactions are not strong enough to support equilibrium, but the mean free path is still much smaller than the typical scales of variation of the distribution function, we can use an approximate description, either with a viscous fluid approximation in the usual context of non-perfect fluids, or using e.g. a tight-coupling expansion [3–8] in the context of cosmic microwave background (CMB) in the early Universe before the electron decoupling

Among the relativistic contexts in which the kinetic theory cannot be avoided, we find *i)* the big-bang nucleosynthesis [2, 9–13] during which the evolution of neutrinos, which are highly relativistic and gradually decouple from other species, needs to be described by a Boltzmann equation; *ii)* supernovae explosions to treat the radiative transfer of neutrinos [14, 15]; *iii)* CMB in the decoupling phase and in the transparent matter dominated era.

In this last case, it happens that the description is not complete if we ignore the polarisation state for the gas of photons since a quadrupolar distribution of velocities induces linear polarisation. Indeed, when coupling to electrons stops to be very efficient around recombination, photons develop a quadrupolar distribution of velocities and this leads to the generation of linear polarisation. In the case of photons, the description of polarization is based on a tensor-valued distribution function $f_{\mu\nu}(t, \mathbf{x}, \mathbf{p})$, which is doubly projected orthogonally to the observer and to the momentum [16, 17]. While the description of polarisation in the kinetic theory is well established for photons, this is not the case for fermions and there are several open questions. First with which type of distribution function can we describe the various polarisation states for a gas of fermions? What are the conditions required to generate or

*Electronic address: christian.fidler@uclouvain.be

†Electronic address: pitrou@iap.fr

to reduce and erase polarisation in a gas of fermions? Under which circumstances polarisation can be ignored and a description based on a simple distribution function is sufficient?

Main features of the formalism

In this article we aim at answering these questions by building the kinetic theory for fermions, allowing for the description of polarisation. We build a meaningful procedure to obtain a classical description out of a quantum system. This is even more crucial for a gas of fermions, which are a set of excitations of an underlying quantum field, since there is no classical version of the field as in the case of bosons, e.g. in the case of photons where the classical field is described by classical electrodynamics. The fundamental step is to consider the number operator $N_{rs}(\mathbf{p}, \mathbf{p}') \equiv a_r^\dagger(\mathbf{p})a_s(\mathbf{p}')$ in the quantum system, where the indices of the creation and annihilation operators a_r^\dagger and a_s refer to helicity. Its expectation value takes a particularly simple form in the case of a homogeneous system since this imposes

$$\langle N_{rs}(\mathbf{p}, \mathbf{p}') \rangle = (2\pi)^3 2p^0 \delta^3(\mathbf{p} - \mathbf{p}') f_{rs}(\mathbf{p}). \quad (1)$$

We can show that the function $f_{rs}(\mathbf{p})$ describes correctly the statistics of momenta and in order to obtain a classical limit we only need to consider that the results obtained in the homogeneous case are valid locally, that is we consider a function $f_{rs}(\mathbf{x}, t, \mathbf{p})$.

An operator in spinor space for fermion particles can then be formed by contraction with the plane-wave solutions of the Dirac equation $u_s(\mathbf{p})$

$$\mathbf{F}(\mathbf{x}, t, \mathbf{p}) \equiv \sum_{rs} u_s(\mathbf{p}) f_{rs}(\mathbf{x}, t, \mathbf{p}) \bar{u}_r(\mathbf{p}), \quad (2)$$

and we show that it is a central object in the computation of the collision term of the Boltzmann equation. The method is slightly different but essentially similar for fermion antiparticles. In full generality an operator in spinor space has 16 degrees of freedom but given its construction based on the number operator, we have only 4 independent degrees of freedom in the operator $\mathbf{F}(\mathbf{x}, t, \mathbf{p})$ (corresponding to the two helicity states and their correlation). We show hereafter that these degrees of freedom are recovered in a covariant and observer independent manner when decomposing the operator (2) as

$$\mathbf{F} = \frac{1}{2} (I - \gamma^5 \gamma^\mu \mathcal{Q}_\mu) (m - \not{p}). \quad (3)$$

One degree of freedom corresponds to the total intensity $I(\mathbf{x}, t, \mathbf{p})$ while the three remaining degrees correspond to the state of polarisation and are covariantly contained in a vector $\mathcal{Q}^\mu(\mathbf{x}, t, \mathbf{p})$ with the transverse property $p_\mu \mathcal{Q}^\mu = 0$.

These degrees of freedom are parallel transported by free streaming and thus allow for a description by a simple Liouville equation. More importantly, they are independent from the observer even though a choice of observer is needed for the construction of helicity states; they are true functions of the phase space. We further show that the collision term for fermion interactions, such as weak interactions, is naturally expressed in these covariant quantities.

Outline

We first summarize the description of fermion quantum fields in § I. We then build in § II the distribution function $\mathbf{F}(\mathbf{x}, t, \mathbf{p})$ which is an operator in spinor space and we show how its covariant components $I(\mathbf{x}, t, \mathbf{p})$ and $\mathcal{Q}^\mu(\mathbf{x}, t, \mathbf{p})$ are extracted. The transformation properties of these quantities are investigated in § III and we show that they are observer independent. Hence they are well suited to describe the free streaming of the distribution function and we derive the Liouville equation that they satisfy in § IV. We then turn to the effect of collisions in § V where we derive the general Boltzmann equation for fermions. This is applied to the specific case of low energy weak interactions in § VI. We detail the form of the collision terms obtained for standard reactions in § VII so as to recover standard results when polarisation can be ignored. The case of vector bosons, including photons, is summarized in appendix A. Finally, technical details are gathered in appendix B for commutation rules of quantum creation and annihilation operators, appendix C for traces of Dirac gamma matrices, and appendix D for two-component spinors and their relation to Dirac spinors.

Notation

An inertial frame is defined by a tetrad field, that is by a timelike vector field e_0 and three spacelike vector fields e_i , together with the associated co-tetrad e^0, e^i . Latin indices such as i, j, \dots indicate spatial components in the tetrad basis. A four-vector is written as V^μ where Greek indices μ, ν, \dots denote components in the tetrad basis. If gravity can be ignored, that is in the context of special relativity, the inertial frame is global. However, when including the effect of gravity in the context of general relativity, the inertial frame is local and one must employ general coordinates whose indices are labelled by α, β, \dots . For a given vector this implies $V^\alpha = V^\mu [e_\mu]^\alpha$.

The momentum vector p^μ will often simply be denoted as p and its spatial components p^i allow to build the spatial momentum $\mathbf{p} = p^i e_i$. More generally, we reserve boldface notation to spatial vectors. The energy associated with the momentum is given by the time component $E = p^0$. When a quantity depends on the spatial momentum, we use indifferently \mathbf{p} or p when no ambiguity can arise. The (special) relativistic (and Lorentz covariant) integration measure is defined as

$$[dp] \equiv \frac{d^3 \mathbf{p}}{(2\pi)^3 2p^0} \quad (4)$$

and its associated (special) relativistic Dirac function is defined accordingly as

$$\underline{\delta}(p - p') = (2\pi)^3 2p^0 \delta^3(\mathbf{p} - \mathbf{p}'), \quad (5)$$

such that $\int [dp] \underline{\delta}(p - p') = 1$. Our metric convention follows the standard notation employed in cosmology. In the tetrad basis, the metric $g_{\mu\nu}$ reduces to the Minkowski metric

$$\eta = \text{diag}(-, +, +, +). \quad (6)$$

The Levi-Civita tensor is fully antisymmetric and in the tetrad basis all its components are deduced from the choice

$$\epsilon_{0123} = -\epsilon^{0123} = 1. \quad (7)$$

We identify the time-like vector of a tetrad e_0 with the velocity u of an observer and its spatial Levi-Civita is obtained from $\epsilon_{ijk} \equiv u^\mu \epsilon_{\mu ijk}$, and it is such that $\epsilon_{123} = \epsilon^{123} = 1$.

I. DESCRIPTION OF FERMIONS IN QUANTUM FIELD THEORY

A. Quantum fields

Let us consider a species of fermions a (e.g. electrons or neutrinos) and the corresponding antiparticle species \bar{a} . The associated quantum field is decomposed as

$$\psi = \sum_{s=\pm\frac{1}{2}} \int [dp] [e^{-ip \cdot x} \bar{a}_s^\dagger(p) v_s(p) + e^{ip \cdot x} a_s(p) u_s(p)]. \quad (8)$$

In this expression, the creation and annihilation operators of the particles and antiparticles satisfy the anti-commutation rules

$$\{a_r(p), a_s^\dagger(p')\} = \delta_{rs}^K \underline{\delta}(p - p'), \quad \{\bar{a}_r(p), \bar{a}_s^\dagger(p')\} = \delta_{rs}^K \underline{\delta}(p - p') \quad (9)$$

with all other anti-commutators vanishing and where δ^K is the Kronecker function. Furthermore, the u_s and v_s are plane-wave solutions of the Dirac equation

$$(i\gamma^\mu \partial_\mu + m)\psi = 0, \quad (10)$$

where the Dirac matrices γ^μ satisfy the Clifford algebra (for more details, see Appendix D 2)

$$\{\gamma^\mu, \gamma^\nu\} = -2g^{\mu\nu} \mathbf{1}. \quad (11)$$

More precisely, the u_s (v_s) are the positive (negative) frequency solutions and satisfy

$$(\not{p} + m)u_s(p) = 0, \quad (\not{p} - m)v_s(p) = 0, \quad (12)$$

with the standard Dirac slashed notation $\not{p} \equiv p_\mu \gamma^\mu$.

B. Helicity basis

The creation and annihilation operators employed to build the field ψ in Eq. (8) refer to a complete set of solutions to the Dirac equation and we are free to choose the basis characterising this Hilbert space. In the following we will employ helicity states $s = \pm\frac{1}{2}$, with left-helical states labelled as $-1/2$ and right-helical ones as $+1/2$. We first introduce the matrix

$$\gamma^5 \equiv \frac{i}{4!} \epsilon_{\mu\nu\rho\lambda} \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\lambda = i\gamma^0 \gamma^1 \gamma^2 \gamma^3, \quad (13)$$

which satisfies $\{\gamma^5, \gamma^\mu\} = 0$ and $(\gamma^5)^2 = \mathbb{1}$, from which the projectors on the left and right chiral parts are built

$$P_- \equiv \frac{1}{2}(\mathbb{1} - \gamma^5), \quad P_+ \equiv \frac{1}{2}(\mathbb{1} + \gamma^5). \quad (14)$$

Let us consider a momentum p with energy $E = p^0$, and whose spatial direction unit vector $\hat{\mathbf{p}} \equiv \mathbf{p}/|\mathbf{p}|$ is decomposed into spherical coordinates as $\hat{p}^i = (\sin(\theta)\sin(\phi), \sin(\theta)\cos(\phi), \cos(\theta))$. We define left- and right-chiral states $u_L = P_- u$ and $u_R = P_+ u$. From Eq. (12), specifying the particle and antiparticle solutions of the Dirac equation, we find

$$(E\gamma^0 - \mathbf{p} \cdot \boldsymbol{\gamma}) u_{s,R} = m u_{s,L} \quad (E\gamma^0 - \mathbf{p} \cdot \boldsymbol{\gamma}) u_{s,L} = m u_{s,R} \quad (15a)$$

$$(E\gamma^0 - \mathbf{p} \cdot \boldsymbol{\gamma}) v_{s,R} = -m v_{s,L} \quad (E\gamma^0 - \mathbf{p} \cdot \boldsymbol{\gamma}) v_{s,L} = -m v_{s,R}, \quad (15b)$$

enforcing a connection between the left- and right-chiral components of particles and antiparticles for massive fermions. Helicity states, describing the spin related to the spatial momentum of the particle, are defined by

$$-\hat{\mathbf{p}} \cdot \boldsymbol{\gamma} \gamma^5 u_s = 2s \gamma^0 u_s \quad -\hat{\mathbf{p}} \cdot \boldsymbol{\gamma} \gamma^5 v_s = -2s \gamma^0 v_s. \quad (16)$$

For helicity states Eqs. (15) thus simplify to

$$(E + 2s|\mathbf{p}|) u_{s,R} = m \gamma^0 u_{s,L} \quad (E - 2s|\mathbf{p}|) u_{s,L} = m \gamma^0 u_{s,R} \quad (17a)$$

$$(E - 2s|\mathbf{p}|) v_{s,R} = -m \gamma^0 v_{s,L} \quad (E + 2s|\mathbf{p}|) v_{s,L} = -m \gamma^0 v_{s,R}. \quad (17b)$$

In the helicity basis for the Dirac matrices Eq. (16) reads

$$\hat{\mathbf{p}} \cdot \boldsymbol{\sigma} u_{s,L/R} = 2s u_{s,L/R} \quad \hat{\mathbf{p}} \cdot \boldsymbol{\sigma} v_{s,L/R} = -2s v_{s,L/R}, \quad (18)$$

and the left- and right-chiral parts are individually solved by the well known chiral 2-spinors of quantum-mechanics

$$\chi_{\frac{1}{2}} = \begin{pmatrix} e^{-i\phi/2} \cos(\theta/2) \\ e^{i\phi/2} \sin(\theta/2) \end{pmatrix} \quad \chi_{-\frac{1}{2}} = \begin{pmatrix} -e^{-i\phi/2} \sin(\theta/2) \\ e^{i\phi/2} \cos(\theta/2) \end{pmatrix}. \quad (19)$$

The full Dirac spinors are then obtained by combining the left- and right-chiral parts under consideration of Eq. (17) and their normalisation

$$u_s = \begin{pmatrix} \sqrt{E + 2s|\mathbf{p}|} \chi_s \\ \sqrt{E - 2s|\mathbf{p}|} \chi_s \end{pmatrix} \quad v_s = \begin{pmatrix} 2s \sqrt{E - 2s|\mathbf{p}|} \chi_{-s} \\ -2s \sqrt{E + 2s|\mathbf{p}|} \chi_{-s} \end{pmatrix}. \quad (20)$$

In Appendix D we detail how these solutions are also expressed with the two-components spinor formalism.

II. QUANTUM DISTRIBUTION FUNCTION

A. Number operator and distribution function

Let us define the number operator

$$N_{rs}(p, p') \equiv a_r^\dagger(p) a_s(p'). \quad (21)$$

If we consider a homogenous system described by the state $|\Psi\rangle$, then its average in this quantum state must be diagonal and of the form

$$\langle \Psi | N_{rs}(p, p') | \Psi \rangle \equiv \delta(p - p') f_{rs}(p). \quad (22)$$

This can be understood when considering the Wigner functional [18]. To that purpose, let us introduce the average and difference of momentum

$$\mathbf{K} \equiv \mathbf{p}' - \mathbf{p} \quad \bar{\mathbf{p}} \equiv \frac{1}{2}(\mathbf{p}' + \mathbf{p}). \quad (23)$$

Then the Wigner transform is defined as

$$\langle N_{rs}(\bar{\mathbf{p}}, \mathbf{x}) \rangle \equiv \int \frac{d^3 \mathbf{K}}{(2\pi)^3 2K^0} \langle \Psi | N_{rs}(p, p') | \Psi \rangle e^{i\mathbf{K} \cdot \mathbf{x}}. \quad (24)$$

Considering a homogeneous system in the sense of the Wigner functional means that $\langle N_{rs}(\bar{\mathbf{p}}, \mathbf{x}) \rangle$ does not depend on \mathbf{x} and in that case it is equal to the function $f_{rs}(\bar{\mathbf{p}})$ defined in Eq. (22).

It is then straightforward to show that this should be identified with the one-particle distribution function of a homogeneous system. Indeed the total occupation operator is obtained from a sum over all possible momenta of the diagonal part as

$$N_{rs} \equiv \int [dp] N_{rs}(p, p). \quad (25)$$

Its average in a given quantum state is

$$\langle \Psi | N_{rs} | \Psi \rangle = \mathcal{V} \int \frac{d^3 \mathbf{p}}{(2\pi)^3} f_{rs}(p), \quad (26)$$

where we introduced the total volume $(2\pi)^3 \delta^3(0) = \mathcal{V}$. This corresponds exactly to the definition of a classical one-particle distribution function. By construction N_{rs} and f_{rs} are Hermitian, that is

$$N_{rs}^*(p, p') = N_{sr}(p', p) \quad \Rightarrow \quad f_{rs}^*(p) = f_{sr}(p). \quad (27)$$

For antiparticles, the construction is equivalent but with the operators of antiparticles instead.

B. Molecular chaos and the Boltzmann approximation

In principle, when considering an interacting system, the one-particle distribution function is not enough to describe it statistically, because n -particle correlation functions are generated by collisions. In order to obtain a description only in terms of a one-particle distribution function, we must assume that the connected part of n -particle functions vanishes and thus that n -particle functions are expressed only in terms of one-particle functions, corresponding to the assumption of molecular chaos. The n -particle number operators for species a are defined as

$$N_{r_1 \dots r_n s_1 \dots s_n}^{(n)} \equiv a_{r_1}^\dagger \dots a_{r_n}^\dagger a_{s_1} \dots a_{s_n}, \quad \Rightarrow \quad N_{r_1 r_2 s_1 s_2}^{(2)} \equiv a_{r_1}^\dagger a_{r_2}^\dagger a_{s_1} a_{s_2}, \quad (28)$$

where for simplicity we omit to write explicitly the momentum dependence of the operators in this section. Their expectation value in a quantum state are then expressed as¹

$$\langle N_{r_1 \dots r_n s_1 \dots s_n}^{(n)} \rangle = \sum_{\sigma \in S_n} (-1)^{n+1} \epsilon(\sigma) \langle N_{r_1 s_{\sigma(1)}}^{(1)} \rangle \dots \langle N_{r_n s_{\sigma(n)}}^{(1)} \rangle \quad \Rightarrow \quad \langle N_{r_1 r_2 s_1 s_2}^{(2)} \rangle = \langle N_{r_1 s_2}^{(1)} \rangle \langle N_{r_2 s_1}^{(1)} \rangle - \langle N_{r_1 s_1}^{(1)} \rangle \langle N_{r_2 s_2}^{(1)} \rangle, \quad (29)$$

where the sum is on the group of permutation S_n and $\epsilon(\sigma)$ is the signature of the permutation. This approximation is exactly similar to the Boltzmann approximation of the BBGKY hierarchy. In practice this assumption of molecular chaos is used to obtain the following property for the expectation in a quantum state of a product of one-particle number operators

$$\langle N_{r_1 s_1}^{(1)} \dots N_{r_n s_n}^{(1)} \rangle = \sum_{\sigma \in S_n} \langle \{N_{r_1 s_{\sigma(1)}}^{(1)}\} \rangle \dots \langle \{N_{r_n s_{\sigma(n)}}^{(1)}\} \rangle \quad \{N_{r_a s_b}^{(1)}\} = \begin{cases} N_{r_a s_b}^{(1)} & a \leq b \\ \widehat{N}_{r_a s_b}^{(1)} & a > b \end{cases}, \quad (30)$$

¹ For bosons we remove all minus signs, that is the factor $(-1)^{n+1} \epsilon(\sigma)$ so we would for instance get $\langle N_{r_1 r_2 s_1 s_2}^{(2)} \rangle = \langle N_{r_1 s_1}^{(1)} \rangle \langle N_{r_2 s_2}^{(1)} \rangle + \langle N_{r_1 s_2}^{(1)} \rangle \langle N_{r_2 s_1}^{(1)} \rangle$.

where \widehat{N}_{rs} is the Pauli blocking operator defined in Eq. (B2). The expectation value of a product of one-particle number operators is the sum of products of expectation values of all possible pairings between creation and annihilation operators. For each pair, if the indices (r_a, s_b) correspond to operators which were initially in the creation-annihilation order, that is with $a \leq b$ (resp. annihilation-creation order, that is $a > b$) we use $N_{r_a s_b}$ (resp. $\widehat{N}_{r_a s_b}$). For instance the expectation value for a product of two one-particle number operators is simply

$$\langle N_{r_1 s_1}^{(1)} N_{r_2 s_2}^{(1)} \rangle = \langle N_{r_1 s_1}^{(1)} \rangle \langle N_{r_2 s_2}^{(1)} \rangle + \langle N_{r_1 s_2}^{(1)} \rangle \langle \widehat{N}_{r_2 s_1}^{(1)} \rangle. \quad (31)$$

Finally, we also assume that species are uncorrelated such that the expectation value for operators of various species is the product of expectation values of the operators of each species. For instance for two species a and b we assume $\langle a_r^\dagger a_s b_p^\dagger b_q \rangle = \langle a_r^\dagger a_s \rangle \langle b_p^\dagger b_q \rangle$.

C. Spinor valued distribution function

In order to extract the information contained in $f_{rs}(p)$ in a covariant form, we build operators in spinor space following

$$F_a^b(p) \equiv \begin{cases} \sum_{rs} f_{rs}(p) u_{s,a}(p) \bar{u}_r^b(p) & \text{particles,} \\ \sum_{rs} f_{rs}(p) v_{r,a}(p) \bar{v}_s^b(p) & \text{antiparticles.} \end{cases} \quad (32)$$

A spinor valued operator has 16 degrees of freedom and we thus need a 16-dimensional basis to decompose operators in spinor space. The set

$$\mathcal{O} \equiv \{\mathbb{1}, \gamma^\mu, \Sigma^{\mu\nu}, \gamma^\mu \gamma^5, \gamma^5\} \quad (33)$$

is a complete basis for the space of operators in spinor space, where we define the matrices

$$\Sigma^{\mu\nu} \equiv \frac{i}{4} [\gamma^\mu, \gamma^\nu], \quad \widetilde{\Sigma}^{\mu\nu} \equiv \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} \Sigma^{\alpha\beta} = i\gamma^5 \Sigma^{\mu\nu}. \quad (34)$$

For the sake of clarity we use a notation where components of operators in spinor space and Dirac spinors are explicit and are denoted by indices of the type $\mathbf{a}, \mathbf{b}, \dots$. For instance the identity operator is δ_a^b and the plane waves solutions are written $u_{s,a}$ and $v_{s,a}$. For any two different operators X and Y (X_a^b and Y_a^b in spinor components) in the set \mathcal{O} , $\text{Tr}[X \cdot Y] = X_a^b Y_b^a = 0$.

Using this orthogonality property, any operator can be decomposed onto this basis with the help of the Fierz identity [19] (see also Eq. G.1.99 of [20] taking into account a factor 2 difference in the definition of $\Sigma^{\mu\nu}$)

$$\delta_a^b \delta_c^d = \frac{1}{4} [\delta_a^d \delta_c^b + (\gamma^5)_a^d (\gamma^5)_c^b - (\gamma^\mu)_a^d (\gamma_\mu)_c^b + (\gamma^\mu \gamma^5)_a^d (\gamma_\mu \gamma^5)_c^b + 2(\Sigma^{\mu\nu})_a^d (\Sigma_{\mu\nu})_c^b] = \sum_{X \in \mathcal{O}} c_X X_a^d X_c^b. \quad (35)$$

The last equality defines the coefficients c_X of the expansion which, by construction, satisfy $c_X = 1/\text{Tr}[X \cdot X]$. Note also that this identity can be used with the matrices $\widetilde{\Sigma}^{\mu\nu}$ instead of $\Sigma^{\mu\nu}$ by employing $(\Sigma^{\mu\nu})_a^d (\Sigma_{\mu\nu})_c^b = -(\widetilde{\Sigma}^{\mu\nu})_a^d (\widetilde{\Sigma}_{\mu\nu})_c^b$, and in the following this proves to be more convenient.

Any operator O in spinor space is decomposed on the basis \mathcal{O} thanks to Eq. (35) as

$$O_a^b = \sum_{c_X \in \mathcal{O}} c_X \text{Tr}[O \cdot X] X_a^b. \quad (36)$$

In particular, any bilinear tensor product of the form $u_r(p) \bar{u}_s(p)$ or $v_r(p) \bar{v}_s(p)$ [with the standard notation $\bar{u}_s = u_s^\dagger \gamma^0$, see Eq. (D20)], is a spinor-space operator and can be decomposed as

$$u_{r,a}(p) \bar{u}_s^b(p) = \sum_{X \in \mathcal{O}} c_X [\bar{u}_s(p) X u_r(p)] X_a^b = \sum_{X \in \mathcal{O}} c_X [\bar{u}_s^c(p) X_c^d u_{r,d}(p)] X_a^b. \quad (37)$$

Hence, we only need to compute the $\bar{u}_s(p) X u_r(p)$ and $\bar{v}_s(p) X v_r(p)$ (these expressions are computed in App. D 7) for all $X \in \mathcal{O}$ to decompose the operator F_a^b , defined in Eq. (32), into

$$F_a^b = \begin{cases} \sum_{X \in \mathcal{O}} c_X \sum_{rs} f_{rs} (\bar{u}_r X u_s) X_a^b & \text{particles,} \\ \sum_{X \in \mathcal{O}} c_X \sum_{rs} f_{sr} (\bar{v}_r X v_s) X_a^b & \text{antiparticles.} \end{cases} \quad (38)$$

Note how the indices are in reverse order for antiparticles (f_{sr} instead of f_{rs}).

1. Covariant components of the spinor valued distribution function

While $f_{rs}(p)$ has only four degrees of freedom, F_a^b has 16 components, and in the decomposition this must be made manifest. To that purpose let us define a set of three unit spatial vectors. For a given observer with four-velocity u^α which we identify with the time-like tetrad vector e_0^α , we can define a spatial momentum \mathbf{p} and its spatial direction $\hat{\mathbf{p}} \equiv \mathbf{p}/|\mathbf{p}|$. In spherical coordinates the momentum direction is given by θ, ϕ and defines a radial unit vector. We then also consider the usual basis in spherical coordinates e_θ and e_ϕ , which are purely spatial unit vectors. In tetrad components these are given by

$$\hat{p}^i = \begin{pmatrix} \cos \phi \sin \theta \\ \sin \phi \sin \theta \\ \cos \theta \end{pmatrix}, \quad e_\theta^i = \begin{pmatrix} \cos \phi \cos \theta \\ \sin \phi \cos \theta \\ -\sin \theta \end{pmatrix}, \quad e_\phi^i = \begin{pmatrix} -\sin \phi \\ \cos \phi \\ 0 \end{pmatrix}. \quad (39)$$

Let us introduce the helicity vector

$$S^\mu(u^\nu, p^\nu) = -\frac{m}{|\mathbf{p}|}u^\mu + \frac{E}{m|\mathbf{p}|}p^\mu, \quad E = -u_\mu p^\mu, \quad |\mathbf{p}| = \sqrt{E^2 - m^2}, \quad (40)$$

which is a unit vector in the direction of the spatial momentum that is transverse to p^μ in the sense $S^\mu p_\mu = 0$ and is thus spacelike. Since the space of vectors orthogonal to p^μ is three-dimensional, the transverse property is not enough to specify the helicity vector and the definition (40) depends explicitly on the observer which is used to define the spatial part of the momentum. When no ambiguity can arise we write simply S^μ . In components the helicity vector is given by

$$S^0 = m^{-1}|\mathbf{p}|, \quad S^i = m^{-1}E\hat{p}^i. \quad (41)$$

Geometrically (see Fig. 1), the helicity vector corresponds to the spatial direction unit vector $\hat{\mathbf{p}}$ boosted in its direction by the same boost needed to obtain p^μ from u^μ .




FIG. 1: Left: In the massive case, p^μ/m and e_θ, e_ϕ, S^μ form an orthonormal basis. Right: In the massless case, k^μ, p^μ are null vectors orthogonal to e_θ, e_ϕ such that $k^\mu p_\mu = -1$. In both cases, the polarisation basis is formed by $\epsilon_\pm^\mu \equiv (e_\theta^\mu \mp i e_\phi^\mu)/\sqrt{2}$

Finally, we define the polarisation basis

$$\epsilon_\pm^\mu \equiv \frac{1}{\sqrt{2}}(e_\theta^\mu \mp i e_\phi^\mu), \quad \Rightarrow \quad \epsilon_\pm^0 = 0, \quad \epsilon_\pm^i = \frac{1}{\sqrt{2}} \begin{pmatrix} \cos(\theta)\cos(\phi) \pm i\sin(\phi) \\ \cos(\theta)\sin(\phi) \mp i\cos(\phi) \\ -\sin(\theta) \end{pmatrix}. \quad (42)$$

The set of vectors S^μ, ϵ_\pm^μ , and p^μ/m constitute an adapted orthonormal basis, with which the expressions for the operators of the type (37) take a simple form. From Eqs. (D45), and noting $m\mathbb{1}$ simply as m , we first recover immediately the known decompositions

$$u_s \bar{u}_s = \frac{1}{2} (\mathbb{1} - 2s\gamma_5 \not{\mathcal{S}}) (-\not{p} + m) = \frac{1}{2} (-\not{p} + m) (\mathbb{1} - 2s\gamma_5 \not{\mathcal{S}}), \quad \sum_s u_s \bar{u}_s = -\not{p} + m \quad (43a)$$

$$v_s \bar{v}_s = \frac{1}{2} (\mathbb{1} - 2s\gamma_5 \not{\mathcal{S}}) (-\not{p} - m) = \frac{1}{2} (-\not{p} - m) (\mathbb{1} - 2s\gamma_5 \not{\mathcal{S}}), \quad \sum_s v_s \bar{v}_s = -\not{p} - m. \quad (43b)$$

Note however that in the massless limit, $mS^\mu \rightarrow p^\mu$ where p^μ is becomes a null vector. The previous decompositions in the massless case take the simpler forms

$$u_s \bar{u}_s = -\frac{1}{2} (\mathbb{1} + 2s\gamma_5) \not{p} = -\frac{1}{2} \not{p} (\mathbb{1} - 2s\gamma_5), \quad \sum_s u_s \bar{u}_s = -\not{p} \quad (44a)$$

$$v_s \bar{v}_s = -\frac{1}{2} (\mathbb{1} - 2s\gamma_5) \not{p} = -\frac{1}{2} \not{p} (\mathbb{1} + 2s\gamma_5), \quad \sum_s v_s \bar{v}_s = -\not{p}. \quad (44b)$$

Additionally we also find corresponding decompositions when the helicities are different, the so-called Bouchiat-Michel formulae [21] (see also App. H.4 of Ref. [20]), and using the polarisation basis (42) these take the simple forms

$$u_s \bar{u}_r = \sqrt{2} p^\mu \epsilon_{r-s}^\nu \tilde{\Sigma}_{\mu\nu} - \frac{m}{\sqrt{2}} \gamma^5 \gamma_\mu \epsilon_{r-s}^\mu = \frac{1}{\sqrt{2}} \gamma^5 \not{\epsilon}_{r-s} (\not{p} - m) \quad \text{if } r \neq s \quad (45a)$$

$$v_s \bar{v}_r = \sqrt{2} p^\mu \epsilon_{s-r}^\nu \tilde{\Sigma}_{\mu\nu} + \frac{m}{\sqrt{2}} \gamma^5 \gamma_\mu \epsilon_{s-r}^\mu = \frac{1}{\sqrt{2}} \gamma^5 \not{\epsilon}_{s-r} (\not{p} + m) \quad \text{if } r \neq s. \quad (45b)$$

We are now in position to decompose the spinor space operator F_a^b . Let us define²

$$I \equiv f_{++} + f_{--}, \quad V = f_{++} - f_{--}, \quad Q_\pm \equiv \sqrt{2} f_{\pm\mp}, \quad (46)$$

together with

$$Q^\mu \equiv Q_+ \epsilon_+^\mu + Q_- \epsilon_-^\mu, \quad \mathcal{Q}^\mu \equiv Q^\mu + VS^\mu. \quad (47)$$

I is the total intensity, V the circular polarisation, and Q^μ is the purely linear polarisation vector. \mathcal{Q}^μ is the total polarisation vector, taking into account both circular and linear polarisation. By construction the total polarisation \mathcal{Q}^μ is transverse to the momentum ($\mathcal{Q}^\mu p_\mu = 0$). The linear polarisation Q^μ is transverse both to the momentum and to the observer velocity u^μ , that is it is a purely spatial vector.

- In the massive case, using Eqs. (43) and (45) in Eq. (38), with again the short notation M for $M\mathbb{1}$, the operator \mathbf{F} is decomposed on the basis of operators \mathcal{O} as

$$\mathbf{F} = \frac{I}{2} (M - \not{p}) - \frac{M}{2} \gamma^5 \gamma_\mu \mathcal{Q}^\mu + p^\mu \tilde{\Sigma}_{\mu\nu} \mathcal{Q}^\nu. \quad (48)$$

We have written this decomposition in a form which is valid for both particles and antiparticles, by introducing the notation

$$M = \begin{cases} m, & \text{particle} \\ -m, & \text{antiparticle.} \end{cases} \quad (49)$$

Using $\gamma^5 \not{p} = \not{p} \gamma^5 \not{Q} = 2p^\mu \tilde{\Sigma}_{\mu\nu} \mathcal{Q}^\nu$ the decomposition can also be written (with the notation I for $I\mathbb{1}$)

$$\mathbf{F} = \frac{1}{2} (I - \gamma^5 \not{Q}) (M - \not{p}) = \frac{1}{2} (M - \not{p}) (I - \gamma^5 \not{Q}). \quad (50)$$

- In the massless limit, we obtain the simpler decomposition

$$\mathbf{F} = -\frac{1}{2} (I + \lambda V \gamma^5) \not{p} + p^\mu \tilde{\Sigma}_{\mu\nu} \mathcal{Q}^\nu, \quad \lambda \equiv \begin{cases} 1 & \text{particles} \\ -1 & \text{antiparticles.} \end{cases} \quad (51)$$

Note that the linear polarisation Q^μ and the circular polarisation V enter separately, and not as a total polarisation vector \mathcal{Q}^μ as is the case in the massive case. Using $\gamma^5 \not{p} = \not{p} \gamma^5 \not{Q} = 2p^\mu \tilde{\Sigma}_{\mu\nu} \mathcal{Q}^\nu$ it can also be rewritten as

$$\mathbf{F} = -\frac{1}{2} (I + \lambda V \gamma^5 - \gamma^5 \not{Q}) \not{p} = -\frac{1}{2} \not{p} (I - \lambda V \gamma^5 - \gamma^5 \not{Q}). \quad (52)$$

The decompositions (51) or (52) match the ones obtained in Refs. [22] [23] which focused on the massless case.

² We use the obvious abuse of notation f_{++} for e.g. $f_{+\frac{1}{2}+\frac{1}{2}}$.

2. Properties of the decomposition in covariant components

- The decomposition of an antiparticle is the same as the one of its particle counterpart, except for the replacement $m \rightarrow -m$. However it is usually admitted that the rule to go from the particle to the antiparticle description is $p \rightarrow -p$. In the decomposition (48), we note that the replacement $m \rightarrow -m$ is equivalent to the replacement $p \rightarrow -p$ up to an overall minus sign. We detail this point in § III E 2.
- In the massless case, the particle and antiparticle operators differ only by the replacement $V \rightarrow -V$.
- All operators X appearing in the decomposition (48) satisfy the property $X^\dagger = \gamma^0 \cdot X \cdot \gamma^0$ or more rigorously $X^\dagger = A \cdot X \cdot A^{-1}$ (see App. D 2 and Eq. G.1.22 of Ref. [20]). Note that this is not the case for the operator γ^5 which does not appear. Hence, the operator \mathbf{F} satisfies the same property

$$\mathbf{F}^\dagger = \gamma^0 \cdot \mathbf{F} \cdot \gamma^0. \quad (53)$$

This property is a consequence of the Hermiticity of the distribution function [see Eq. (27)].

- In the decomposition of the massless case (51), we remark that any extra term proportional to p^μ can be added to Q^μ without altering the decomposition due to the antisymmetry of $\tilde{\Sigma}_{\mu\nu}$, and without altering the transverse property $p^\mu Q_\mu = 0$. Indeed Q^μ has three degrees of freedom since it is built from V, Q_+, Q_- and this is reflected by the transverse property $Q^\mu p_\mu = 0$, but Q^μ contains only the two degrees of freedom corresponding to Q_-, Q_+ and this leads to this ambiguity. This is identical to the gauge freedom of the massless photon which is remaining even in the Lorentz gauge $k_\mu A^\mu = 0$ since it has only two physical degrees of freedom and not three as in the massive case (Proca theory). The solution to this problem is exactly the same, and by construction of Q^μ we have demanded an extra condition which is observer dependent, $Q_\mu u^\mu = 0$. This means that for this observer the linear polarisation vector Q^μ must be purely spatial. This condition is analogous to the observer dependent Coulomb gauge choice ($A^0 = 0$) required to fully fix the gauge potential. However, the physical results do not depend on this Coulomb-type gauge choice since it does not alter the decomposition (51). To summarize, linear polarisation in the massless case is the coset

$$[Q^\mu] \equiv \{Q^\mu + \alpha p^\mu, \alpha \in \mathbb{R}\} \quad (54)$$

whose preferred representative element for a given observer is the one defined in Eq. (47), which is the only one purely spatial for that observer.

- Except in the massless case, the total polarisation vector is a spin-1 representation of $SO(3) \simeq SU(2)$ and the intensity is a spin-0 representation. Indeed, when forming the number operator (21), and thus f_{rs} , we are building the tensor product of spin-1/2 representations and what we have achieved is a decomposition of the reducible representation $\mathbf{2} \otimes \mathbf{2}$ in irreducible components $\mathbf{3} \oplus \mathbf{1}$, where we have denoted $\mathbf{1}, \mathbf{2}, \mathbf{3}$ the spin-0, 1/2, 1 representations of $SU(2)$. In the massless case, the little group of the Lorentz group (see Ref. [24]) is not $SO(3)$ but $SO(2) \simeq U(1)$. Hence the decomposition in irreducible representations is of the form $\mathbf{2}_1 \oplus \mathbf{1} \oplus \mathbf{1}$ where the purely linear polarisation is in the spin-1 representation of $SO(2)$ (noted $\mathbf{2}_1$) and circular polarisation is in the representation $\mathbf{1}$. See Appendix A for the decomposition in irreducible components of the distribution function for massless vector bosons which obeys the same logic.

3. Extraction of covariant components

The covariant components can be obtained by multiplying \mathbf{F} with the appropriate operator and taking the trace, using the Fierz identity (35). In the massive case we find

$$I = \frac{1}{2M} \text{Tr}[\mathbf{F}] = -\frac{1}{2m^2} \text{Tr}[\mathbf{F} \cdot \not{p}] = \frac{1}{4m^2} \text{Tr}[\mathbf{F} \cdot (M - \not{p})] \quad (55a)$$

$$Q^\mu = \frac{1}{2M} \text{Tr}[\mathbf{F} \cdot \gamma^\mu \gamma^5] = -\frac{1}{2m^2} \text{Tr}[\mathbf{F} \cdot \gamma^\mu \gamma^5 \not{p}] = \frac{1}{4m^2} \text{Tr}[\mathbf{F} \cdot \gamma^\mu \gamma^5 (M - \not{p})] = \frac{1}{4m^2} \text{Tr}[\mathbf{F} \cdot (M - \not{p}) \gamma^\mu \gamma^5]. \quad (55b)$$

In the last equality we have used $\{\not{p}, \gamma^\mu \gamma^5\} = -2\gamma^5 p^\mu$ and $\text{Tr}[\mathbf{F} \cdot \gamma^5] = 0$.

In the massless case, we must first introduce a future directed null vector k^μ (see Fig. 1) such that

$$k^\mu p_\mu = -1, \quad k_\mu \epsilon_\pm^\mu = 0, \quad (56)$$

that is which lies in the plane spanned by the observer four-velocity u^μ and p^μ . The covariant parts are then obtained from

$$I = -\frac{1}{2}\text{Tr}[\mathbf{F} \cdot \mathbb{k}] \quad (57a)$$

$$\lambda V = -\frac{1}{2}\text{Tr}[\mathbf{F} \cdot \mathbb{k}\gamma^5] \quad (57b)$$

$$Q^\mu = -\frac{1}{2}\text{Tr}[\mathbf{F} \cdot \gamma^\mu \gamma^5 \mathbb{k}] = -\frac{1}{2}\text{Tr}[\mathbf{F} \cdot \mathbb{k}\gamma^\mu \gamma^5]. \quad (57c)$$

Note that given the decomposition in the massless case (51), the linear polarisation extracted with the previous expressions is automatically transverse to the observer four-velocity u^μ and to the momentum p^μ . That is, introducing a symmetric screen projector $\mathcal{H}_{\mu\nu}$ such that $\mathcal{H}_\mu^\sigma \mathcal{H}_{\sigma\nu} = \mathcal{H}_{\mu\nu}$ and $\mathcal{H}_{\mu\nu} p^\nu = \mathcal{H}_{\mu\nu} u^\nu = 0$, which is built as

$$\mathcal{H}^\mu{}_\nu \equiv \delta^\mu_\nu + p^\mu k_\nu + k^\mu p_\nu = \epsilon_+^\mu \epsilon_{-\nu} + \epsilon_-^\mu \epsilon_{+\nu}, \quad (58)$$

it satisfies automatically $\mathcal{H}^\mu{}_\nu Q^\nu = Q^\mu$.

III. TRANSFORMATION PROPERTIES AND OBSERVER INDEPENDENCE

A. The definition of observer independence

In order to discuss the observer dependence of our quantities, we first have to define the observer. So far we have performed the computations in special relativity in a Minkowski space-time, that is with a global inertial frame. However, we intend to embed this space-time into the general relativistic manifold and define our observer by a fully relativistic trajectory.

From the equivalence principle, the Minkowski space becomes in the context of general relativity the tangent space, and it is connected to the manifold by the tetrads $[e_\mu]^\alpha$ carrying a Lorentz-index inside the brackets and a general relativistic index on the outside. The tetrads are not unique and at each point in space-time a group of tetrads exist, corresponding to the Lorentz symmetry of special relativity.

In the following, we will choose the unique tetrad corresponding to the observer velocity and its spatial orientation (such that the observer velocity in tetrad indices is purely temporal, i.e. the observer is not moving in its own frame). The corresponding local Minkowski space-time is thus identified directly with our relativistic observer and Lorentz transformations of the Minkowski space-time represent a change of observer. We split the general relativistic 4-momentum P^α ³ into a covariant energy and spatial momentum according to

$$P^\alpha = p^\mu [e_\mu]^\alpha = E[e_0]^\alpha + p^i [e_i]^\alpha, \quad (59)$$

where we recall that we have identified the observer velocity with the temporal tetrad $u^\alpha = [e_0]^\alpha$. E and p^i are by construction invariant under general coordinate transformations, but not under a change of the observer. Each observer will measure a different energy or spatial momentum corresponding to its own velocity and the orientation of the spatial vectors in its tetrad.

We will call a quantity observer independent if under a change of observer (equivalent to a local Lorentz transformation) it transforms in a given representation of the Lorentz group. This implies that the same object will be described differently by two observers, but this difference is entirely related to the local tetrad basis of the observer. One example is the local 4-momentum p^μ , transforming as a 4-vector, or the scalar mass of particles m .

On the other hand a quantity is observer-dependent if it does not follow the usual Lorentz transformations. This is for example the case for the local observer velocity $u^\mu = \delta_0^\mu$, which by definition takes this value for each observer and thus does not transform as a vector. In particular this is also the case for the helicity vector S^μ , which directly depends on the observer-dependant u^μ . The choice of observer therefore affects the helicity states and the same quantum state may be described both as left- or right-handed depending on the observer. We conclude that the distribution functions f_{rs} , the energy E and the three momentum p^i are observer-dependent.

³ For clarity we use the notation p^μ for the components of the momentum in the local tetrad basis e^μ and P^α for the components of the same momentum in a general coordinates basis, that is we use both a different symbol and a different type of index.

B. Observer independence of F_a^b

The field ψ is observer independent by construction and does transform in the usual spinor-representation. We can build ψ for two different observers connected by the passive Lorentz transformation Λ , which thus acts actively on the coordinates $\tilde{x} = \Lambda x$, \tilde{x} being the coordinates of a given point for the new observer and x the coordinates of the same point for the original one. In addition to the coordinates change, each observer also defines his own Hilbert space basis and therefore describes quantum states and operators differently. We take this into account via the unitary transformation $U(\Lambda)$ and we find the usual transformation rules for a spinor valued operator

$$\tilde{\psi}_a(\tilde{x}) = D_a^b(\Lambda)U(\Lambda)\psi_b(x)U(\Lambda^{-1}), \quad (60)$$

where $D_a^b(\Lambda)$ is the representation of the Lorentz transformation on the spinor space whose explicit expression is given in Appendix D3. We directly build the fields in both coordinate systems, with both observers defining the momentum \mathbf{p} and the helicity s based on their own coordinates

$$\psi_a(x) = \sum_{s=\pm\frac{1}{2}} \int [dp] [e^{-ip \cdot x} \bar{a}_s^\dagger(p)v_{s,a}(p) + e^{ip \cdot x} a_s(p)u_{s,a}(p)] \quad (61)$$

$$\begin{aligned} \tilde{\psi}_a(\tilde{x}) &= \sum_{s=\pm\frac{1}{2}} \int [d\tilde{p}] [e^{-i\tilde{p} \cdot \tilde{x}} \bar{a}_s^\dagger(\tilde{p})v_{s,a}(\tilde{p}) + e^{i\tilde{p} \cdot \tilde{x}} a_s(\tilde{p})u_{s,a}(\tilde{p})] \\ &= \sum_{s=\pm\frac{1}{2}} \int [dp] [e^{-ip \cdot x} \bar{a}_s^\dagger(\Lambda p)v_{s,a}(\Lambda p) + e^{ip \cdot x} a_s(\Lambda p)u_{s,a}(\Lambda p)], \end{aligned} \quad (62)$$

where in the last line we have defined $\tilde{p} = \Lambda p$ and used that the scalar product $p \cdot x$ is independent of the coordinates and that the integration element $[dp]$ is Lorentz invariant. Combined with Eq. (60) this implies that

$$\sum_s a_s(\Lambda p)u_{s,a}(\Lambda p) = \sum_s D_a^b(\Lambda)U(\Lambda)a_s(p)U(\Lambda^{-1})u_{s,b}(p) \quad (63a)$$

$$\sum_s \bar{a}_s^\dagger(\Lambda p)v_{s,a}(\Lambda p) = \sum_s D_a^b(\Lambda)U(\Lambda)\bar{a}_s^\dagger(p)U(\Lambda^{-1})v_{s,b}(p). \quad (63b)$$

These equations mean, that even though the helicity states (defined by $a_s(p)$ and $a_s(\Lambda p)$) are observer dependent, after summing them with the corresponding spinors $u_{s,a}$ the combination is observer independent and transforms as a regular Dirac spinor valued operator.

The conjugated field $\bar{\psi}^a$ provides the analogous relations

$$\sum_s a_s^\dagger(\Lambda p)\bar{u}_s^a(\Lambda p) = \sum_s U(\Lambda)a_s^\dagger(p)U(\Lambda^{-1})\bar{u}_s^b(p)D_b^a(\Lambda^{-1}) \quad (64a)$$

$$\sum_s \bar{a}_s(\Lambda p)\bar{v}_s^a(\Lambda p) = \sum_s U(\Lambda)\bar{a}_s(p)U(\Lambda^{-1})\bar{v}_s^b(p)D_b^a(\Lambda^{-1}), \quad (64b)$$

or they can be deduced using the property (D19).

We deduce that a change of coordinates is equivalent to a change in the Hilbert basis

$$\begin{aligned} \psi_a(\Lambda x) &= \sum_{s=\pm\frac{1}{2}} \int [dp] [e^{-ip \cdot \Lambda x} \bar{a}_s^\dagger(p)v_{s,a}(p) + e^{ip \cdot \Lambda x} a_s(p)u_{s,a}(p)] \\ &= \sum_{s=\pm\frac{1}{2}} \int [d\tilde{p}] [e^{-i\tilde{p} \cdot \tilde{x}} \bar{a}_s^\dagger(\tilde{p})v_{s,a}(\tilde{p}) + e^{i\tilde{p} \cdot \tilde{x}} a_s(\tilde{p})u_{s,a}(\tilde{p})] \\ &= D_a^b(\Lambda)U(\Lambda) \sum_{s=\pm\frac{1}{2}} \int [d\tilde{p}] [e^{-i\tilde{p} \cdot \tilde{x}} \bar{a}_s^\dagger(\tilde{p})v_{s,b}(\tilde{p}) + e^{i\tilde{p} \cdot \tilde{x}} a_s(\tilde{p})u_{s,b}(\tilde{p})] U^{-1}(\Lambda) \\ &= D_a^b(\Lambda)U(\Lambda)\psi_b(x)U^{-1}(\Lambda). \end{aligned} \quad (65)$$

So we find that the field must satisfy

$$U(\Lambda)\psi_a(x)U(\Lambda^{-1}) = D_a^b(\Lambda^{-1})\psi_b(\Lambda x). \quad (66)$$

We have seen that the distribution functions f_{rs} are observer dependent. However, using the above equations, the tensor F_a^b built from the distribution functions turns out to be entirely observer independent, for both particles and antiparticles. Indeed, under the change of observer, the quantum state $|\Psi\rangle$ seen by the initial observer is seen as $|\widetilde{\Psi}\rangle = U(\Lambda)|\Psi\rangle$ for the new observer, and a momentum p seen by the old observer is now seen as $\tilde{p} \equiv \Lambda p$ by the new observer. From Eqs. (63) and (64) the spinor valued operator \tilde{F}_a^b defined by the new observer is related to the former by

$$\tilde{F}_a^b(\tilde{p}) = D_a^{a'}(\Lambda)F_{a'b'}(p)D_{b'}^b(\Lambda^{-1}). \quad (67)$$

This is the expected transformation rule for spinor valued tensors in momentum space, and according to the definition of § III A, this proves that F_a^b is observer independent.

C. Transformation properties of the covariant components

The Dirac matrices satisfy the transformation rules

$$D(\Lambda)\gamma^\mu D(\Lambda^{-1}) = \gamma^\nu \Lambda_\nu^\mu \quad \Rightarrow \quad D(\Lambda^{-1})\gamma^\mu D(\Lambda) = \Lambda^\mu_\nu \gamma^\nu. \quad (68)$$

For the proper orthochronous Lorentz group $SO^+(1,3)$ this implies that γ^5 is invariant. Combining this property with the transformation rule (67) and the decomposition (48) we deduce that its covariant components transform under a coordinate transformation $\Lambda \in SO^+(1,3)$ as

$$\tilde{I}(\tilde{p}) = I(p), \quad \tilde{Q}^\mu(\tilde{p}) = \Lambda^\mu_\nu Q^\nu(p). \quad (69)$$

This means that they transform exactly as a scalar and vector fields and according to the discussion of § III A, they are therefore observer independent.

The same analysis can be carried in the massless case using the decomposition (50) and we deduce that the covariant components transform as

$$\tilde{I}(\tilde{p}) = I(p), \quad \tilde{V}(\tilde{p}) = V(p), \quad \tilde{Q}^\mu(\tilde{p}) = \tilde{\mathcal{H}}^\mu_\sigma \Lambda^\sigma_\nu Q^\nu(p). \quad (70)$$

The screen projector [see Def. (58)] associated with the new observer and the new momentum, $\tilde{\mathcal{H}}^\mu_\sigma$, ensures that the linear polarisation remains spatial for the new observer according to the discussion of § II C 2. Hence in the massless case, the linear polarisation part is not strictly observer independent, but since this dependence introduced by the screen projector is there only as the result of a choice to remove a non physical degree of freedom, we can still conclude that in that sense the covariant components are observer independent. More rigorously, the coset of linear polarisation [see definition (54)] is observer independent and only the special choice of its representative element is observer dependent and we should rather write the transformation rule of linear polarisation cosets which is $[\tilde{Q}^\mu(\tilde{p})] = [\Lambda^\mu_\nu Q^\nu(p)]$, for which the observer independence is manifest.

D. Discussion

The scalar I describes the total intensity of the field and is expected to be observer independent as the local number of particles is identical for each observer. The information of the polarisation is contained in the vector Q^μ independent of the observer.

On the other hand the parameters V and Q_\pm , describing individually the circular and linear polarisations are not observer independent. The circular polarisation V , for example, changes if the observer is boosted and overtakes the measured particle. We have defined

$$Q^\mu = Q^\mu + VS^\mu, \quad (71)$$

where Q^μ combines multiple observer dependant quantities into one observer independent vector. In the example of the observer overtaking a particle we change all left-helical $-\frac{1}{2}$ states into right-helical $+\frac{1}{2}$ states. This means that the boosted observer will find $\tilde{V} = -V$. At the same time the vector S^μ is also observer dependent and the new observer will define the spatial momentum of the particles with the opposite sign. Therefore the combination VS^μ is invariant under this boost. At the same time the off-diagonal distributions are swapped: $\tilde{f}_{+-} = f_{-+}$. However these

are combined with the polarisation vectors ϵ_{\pm} to form Q^{μ} , which are also interchanged for the new observer, leading to Q^{μ} being invariant under this change.

In a more general case Q^{μ} and V cannot be disentangled in an observer independent manner and there always exists a subset of observers, all related by boosts along the momentum direction and rotations around the momentum direction, that will perceive the field to be entirely circularly polarised without any linear polarisation. For this reason we will work with the observer independent polarisation vector Q^{μ} and only refer to the circular and linear polarisations when we have specified an observer.

In the massless case the situation simplifies. It is no longer possible to overtake the particles as they move at the speed of light in any coordinate system. This leads to both, the circular and linear polarisations V and Q^{μ} [more rigorously the coset of linear polarisation (54)] to be individually observer independent.

E. Discrete transformations

We can repeat the analysis with discrete transformations. We are most notably interested in parity and charge conjugations which are needed to discuss weak interactions.

1. Parity transformation

For parity transformation, the unitary operator \mathcal{P} implements the transformation through

$$\mathcal{P}a_s(\mathbf{p})\mathcal{P} = ia_{-s}(-\mathbf{p}), \quad \mathcal{P}b_s(\mathbf{p})\mathcal{P} = -ib_{-s}(-\mathbf{p}). \quad (72)$$

If we consider the parity transformed quantum system $\mathcal{P}|\Psi\rangle$ then the distribution function $\tilde{f}_{rs}(\mathbf{p})$ built from it satisfies

$$\tilde{f}_{rs}(\mathbf{p}) = f_{-r-s}(-\mathbf{p}), \quad (73)$$

that is all helicities and momenta are reversed. If we now consider the spinor valued operator $\tilde{\mathbf{F}}$ which is built from \tilde{f}_{rs} , and using the property

$$u_{-s}(-\mathbf{p}) = -i\gamma^0 u_s(\mathbf{p}), \quad v_{-s}(-\mathbf{p}) = -i\gamma^0 v_s(\mathbf{p}), \quad (74)$$

we find that it is related to the original operator as

$$\tilde{\mathbf{F}}(-\mathbf{p}) = \gamma^0 \cdot \mathbf{F}(\mathbf{p}) \cdot \gamma^0 = \mathbf{F}^{\dagger}(\mathbf{p}). \quad (75)$$

which is nothing but Eq. (67) applied for a discrete parity transformation with $D(\mathcal{P}) = \gamma^0$. Since

$$\gamma^0 \cdot \gamma^{\mu} \cdot \gamma^0 = (-1)^{\mu} \gamma^{\mu} \quad (76a)$$

$$\gamma^0 \cdot \gamma^{\mu} \gamma^5 \cdot \gamma^0 = -(-1)^{\mu} \gamma^{\mu} \gamma^5, \quad (76b)$$

where we used the compact notation [25]

$$(-1)^{\mu} = \begin{cases} 1 & \mu = 0 \\ -1 & \mu = 1, 2, 3, \end{cases} \quad (77)$$

then we deduce that

$$\tilde{Q}^0(-\mathbf{p}) = -Q^0(\mathbf{p}) \quad \tilde{Q}^i(-\mathbf{p}) = Q^i(\mathbf{p}). \quad (78)$$

The spatial part of the polarisation is unchanged and transforms like an axial vector. The time component is reverted just to ensure that the transverse property $p^{\mu}Q_{\mu}$ still holds after a parity transformation.

2. Charge conjugation

Let us now consider charge conjugation transformations. They are implemented through the unitary operator \mathcal{C} as

$$\mathcal{C}a_s(\mathbf{p})\mathcal{C} = b_s(\mathbf{p}), \quad \mathcal{C}b_s(\mathbf{p})\mathcal{C} = a_s(\mathbf{p}). \quad (79)$$

Hence, the distribution function of a charged conjugated system $\mathcal{C}|\Psi\rangle$ is related to the distribution function of the original state $|\Psi\rangle$ by

$$\tilde{f}_{rs}(\mathbf{p}) = f_{rs}(\mathbf{p}). \quad (80)$$

The corresponding spinor valued operators can be related using

$$u_{s,a}(\mathbf{p}) = C_{ab} \bar{v}_s^b(\mathbf{p}) \quad \bar{u}_s^a(\mathbf{p}) = C^{-1ab} v_{s,b}(\mathbf{p}), \quad (81)$$

where in the chiral representations the components of C_{ab} are antisymmetric and such that

$$C = (i\gamma^0\gamma^2) \quad C^{-1} = -(i\gamma^0\gamma^2). \quad (82)$$

From (80) and (81) and the definitions (38) of the spinor valued operators we get

$$\tilde{F}_a^b(p) = -C_{ac} F_b^c(p) C^{-1ab} \quad \Rightarrow \quad \tilde{\mathbf{F}}(p) = -C \mathbf{F}^T(p) C^{-1}. \quad (83)$$

Since charge conjugation is not a special case of Lorentz transformations, it does not take the form (67). Using the properties (see G.1.24 of [20])

$$C\Gamma^T C^{-1} = \eta_\Gamma^C \Gamma, \quad \eta_\Gamma^C = \begin{cases} +1, & \Gamma = \mathbf{1}, \gamma^5, \gamma^\mu \gamma^5 \\ -1, & \Gamma = \gamma^\mu, \Sigma^{\mu\nu} \end{cases} \quad (84)$$

we deduce that $\tilde{\mathbf{F}}(p)$ is deduced from $\mathbf{F}(p)$ by a global sign change and the replacement $p^\mu \rightarrow -p^\mu$, which is also equivalent to $m \rightarrow -m$, as already noticed in § II C.

IV. KINETIC THEORY FOR FERMIONS IN CURVED SPACETIME

A. Liouville equation

In order to derive a Liouville equation in curved space-time which describes the evolution of the covariant components, we must distinguish between the massive and the massless case.

1. Massive fermions

In the previous sections we have shown that I and \mathcal{Q} are observer independent. In addition, in the local Minkowski frame, they are also parallel transported in the absence of collisions. The helicity of particles does not change in free propagation and, considering that the momentum p^μ is conserved, the vectors ϵ_\pm^μ and S^μ used to build the quantities I and \mathcal{Q} remain unchanged. Hence, in the local Minkowski space we obtain the equations of motion

$$\frac{dI}{dt} = 0, \quad \frac{d\mathcal{Q}^\mu}{dt} = 0. \quad (85)$$

From the point of view of general relativity, these equations are only valid locally and neglect entirely the impact of the relativistic space-time. The intensity I describes the total number of particles. The conservation of I in the absence of collisions in Eq. (85) is equivalent to mass or particle number conservation. The geometrical impact of relativity does not change the number of particles and we may generalise the equation of motion by requiring the conservation of I along a full geodesic

$$\frac{DI}{D\lambda} = 0, \quad (86)$$

where $\frac{D}{D\lambda}$ is the derivative along the particle trajectory parameterized by λ .

The vector \mathcal{Q} is parallel transported in the local space-time and describes polarisation of particles in an observer-independent way. Again, the geometrical nature of general relativity does not change the polarisation of particles and in general relativity we require that \mathcal{Q} is parallel transported along the non-trivial trajectory of the particles.

Using the observer-independence, we are able to uniquely define the vector \mathcal{Q} on our full space-time by employing the tetrads

$$\mathcal{Q}^\alpha = \mathcal{Q}^\mu [e_\mu]^\alpha, \quad (87)$$

where we remind that the index μ is a tetrad component index, but the index α is a general coordinate index. Assuming parallel transport, we obtain the equation of motion

$$\frac{D\mathcal{Q}^\alpha}{D\lambda} = 0. \quad (88)$$

Note that \mathcal{Q} is by definition orthogonal to the momentum P . This property is automatically conserved in the relativistic evolution as both, P and \mathcal{Q} are parallel-transported along the geodesic of a free particle.

The variation of coordinates along the trajectory is given by $dx^\alpha/d\lambda = P^\alpha$. The derivatives along the trajectory can be expressed in terms of the time-coordinate $\tau = x^0$ using $d\tau/d\lambda = P^0$, and using the dependencies $I = I(\tau, \mathbf{x}, \mathbf{p})$ we find

$$\frac{\partial I(\tau, \mathbf{x}, \mathbf{p})}{\partial \tau} + \frac{P^i}{P^0} \frac{\partial I(\tau, \mathbf{x}, \mathbf{p})}{\partial x^i} + \frac{dp^i}{d\tau} \frac{\partial I(\tau, \mathbf{x}, \mathbf{p})}{\partial p^i} = 0, \quad (89)$$

where we stress that the momentum appearing in the intensity $I(\tau, \mathbf{x}, \mathbf{p})$ is the local 3-momentum \mathbf{p} whose components are expressed in the local tetrad basis of the observer. For the polarisation vector we obtain

$$\frac{\partial \mathcal{Q}^\alpha(\tau, \mathbf{x}, \mathbf{p})}{\partial \tau} + \frac{P^i}{P^0} \frac{\partial \mathcal{Q}^\alpha(\tau, \mathbf{x}, \mathbf{p})}{\partial x^i} + \frac{dp^i}{d\tau} \frac{\partial \mathcal{Q}^\alpha(\tau, \mathbf{x}, \mathbf{p})}{\partial p^i} + \Gamma_{\gamma\beta}^\alpha \frac{P^\gamma}{P^0} \mathcal{Q}^\beta(\tau, \mathbf{x}, \mathbf{p}) = 0, \quad (90)$$

where the $\Gamma_{\gamma\beta}^\alpha$ are the Christoffel symbols associated with the metric. These are the usual relativistic Boltzmann equations, with the second term describing the variation of the distribution function due to particle propagation and the third term describing the impact of a change of the local momentum (e.g. due to redshifting related to the expansion of the Universe). For the propagation of the polarisation vector, we find an additional geometrical term from the non-trivial parallel transport of vectors in curved spaces.

Finally, the change of the local momentum $dp^i/d\tau$ is obtained from the geodesic equation

$$\frac{DP^\alpha}{D\lambda} = \frac{dP^\alpha}{d\lambda} + \Gamma_{\gamma\beta}^\alpha P^\beta P^\gamma = 0 \quad \Rightarrow \quad \frac{dp^\mu}{d\tau} = P^\alpha \frac{d[e^\mu]_\alpha}{d\tau} - [e^\mu]_\alpha \Gamma_{\gamma\beta}^\alpha \frac{P^\beta P^\gamma}{P^0}. \quad (91)$$

2. Massless fermions

In the massless case, linear polarisation and circular polarisation must be considered separately. Circular polarisation V is transported exactly like the intensity I in Eqs. (86) and (89) because the direction of the helicity vector is identical to the momentum and therefore parallel-transported. However the linear polarisation vector (considered in general coordinates with $Q^\alpha = \mathcal{Q}^\mu [e_\mu]^\alpha$) cannot be parallel transported because it is transverse to both the momentum P and the observer velocity identified with e_0 , and the latter is not parallel transported. In the process of free streaming, any variation of Q^α in the direction of P^α is not physical as explained at the end of § (II C 1). Hence this unphysical degree of freedom must be eliminated by an appropriate projection so as to obtain an unambiguous equation for parallel transport. To that purpose, we use the screen projector (58) in general coordinates. By construction $\mathcal{H}^\alpha_\beta Q^\beta = Q^\alpha$ and linear polarisation is parallel transported *as much as the double transverse property allows* as

$$\mathcal{H}^\beta_\alpha \frac{DQ^\alpha}{D\lambda} = 0. \quad (92)$$

The transport of linear polarisation in the massless case is the same as the transport of the full polarisation vector in the massive case [Eqs. (88) and (90)], up to an additional screen projection which ensures that the double transverse property holds. This is similar to the parallel transport of linear polarisation for photons [16, 17, 26, 27] which in that case is described by a doubly projected tensor (see App. A for more details about the statistical description of vector bosons).

B. Angular decomposition and multipoles

The intensity $I(\mathbf{p})$ is easily decomposed into spherical harmonics. Indeed, once an observer choice is made, that is its four-velocity u^α is identified with the time-like vector of the tetrad $[e_0]^\alpha$, we can define the spatial momentum \mathbf{p} and its direction unit vector $\hat{\mathbf{p}}$. We then perform the usual spherical harmonics decomposition

$$I(\mathbf{p}) = \sum_{\ell m} I_{\ell m}(|\mathbf{p}|) Y_{\ell m}(\hat{\mathbf{p}}). \quad (93)$$

Alternatively one could utilise a decomposition based on symmetric trace-free tensors which is equivalent [17, 28].

For the polarisation vector Q^μ we remind ourselves of its decomposition as

$$Q^\mu(\mathbf{p}) = Q_+(\mathbf{p})\epsilon_+^\mu(\mathbf{p}) + Q_-(\mathbf{p})\epsilon_-^\mu(\mathbf{p}) + V(\mathbf{p})S^\mu(\mathbf{p}). \quad (94)$$

For the angular decomposition we have to pay attention to the transformation properties when performing a spatial rotation of the coordinate system around the direction of $\hat{\mathbf{p}}$. The ordinary spherical harmonics, when evaluated at $\hat{\mathbf{p}}$ do not transform under this rotation and are thus not suitable to decompose objects which have a non-trivial transformation under this rotation. The polarisation vector Q transforms as an ordinary 4-vector (we have shown that it is observer-independent). However, this is not the case for the observer-dependent vectors and distribution functions used to build Q . The vector in direction of the spatial momentum S^μ is invariant under this particular rotation as it points in the direction $\hat{\mathbf{p}}$. Employing the observer-independence of Q we therefore conclude that V must be invariant under this rotation and may be decomposed into ordinary spherical harmonics. The polarisation vectors $\epsilon_\pm(\hat{\mathbf{p}})$ however transform with an additional spin ∓ 1 complex rotation. To generate an observer-independent Q the corresponding Q_\pm must transform with the opposite spin and they are decomposed into spin-weighted spherical harmonics $Y_{\ell m}^s$ [29] as

$$Q_+(\mathbf{p}) \equiv \sum_{\ell m} Q_{\ell m}^+(|\mathbf{p}|) Y_{\ell m}^+(\hat{\mathbf{p}}), \quad Q_-(\mathbf{p}) \equiv \sum_{\ell m} Q_{\ell m}^-(|\mathbf{p}|) Y_{\ell m}^-(\hat{\mathbf{p}}). \quad (95)$$

Note that this discussion only concerns the observer dependence under a specific spatial rotation and that due to the definition of helicity an additional dependence mixing V and Q_\pm exists for more general rotations and boosts.

E and B modes multipoles can be defined from $Q_{\ell m}^\pm \equiv \mp(E_{\ell m} \pm iB_{\ell m})$. Equivalently since $Q(\mathbf{p})$ is a vector field on the unit sphere in momentum space, it can be decomposed as the gradient and the curl of two scalar functions as

$$Q_i(\mathbf{p}) = D_i \mathcal{E}(\mathbf{p}) + \hat{p}^j \epsilon_{ji}{}^k D_k \mathcal{B}(\mathbf{p}), \quad (96)$$

where D_i is the covariant derivative on the unit sphere and $\hat{p}^j \epsilon_{ji}{}^k$ is the Levi-Civita tensor on the unit sphere. Decomposing the scalar functions \mathcal{E} and \mathcal{B} in multipoles $\mathcal{E}_{\ell m}$ and $\mathcal{B}_{\ell m}$ as in the expansion (93) and using [30]

$$D^i Y_{\ell m} = \sqrt{\frac{\ell(\ell+1)}{2}} (-Y_{\ell m}^+ \epsilon_+^i + Y_{\ell m}^- \epsilon_-^i) \quad (97)$$

the two possible definitions for the E and B modes multipoles are related by $E_{\ell m} = \sqrt{\ell(\ell+1)/2} \mathcal{E}_{\ell m}$ and $B_{\ell m} = \sqrt{\ell(\ell+1)/2} \mathcal{B}_{\ell m}$. Again a similar expansion can be obtained by using symmetric trace-free tensors to expand the scalar functions \mathcal{E} and \mathcal{B} directly in Eq. (96).

V. THE BOLTZMANN EQUATION

A. Time evolution and collisions

So far we have discussed the free propagation of fermions. When considering collisions, we will employ a separation of scales. We assume that the relativistic evolution is dominant on macroscopic scales, while individual collisions act on microscopic scales. We therefore may compute the collision term in the local tangent space corresponding to special relativity. Then averaging over the local Minkowski space-time of the observer we will provide an effective collision term for the relativistic evolution of the distribution functions.

We therefore introduce three relevant scales. First the microscopic scale of individual interactions, typically the Compton timescale of interacting particles. Then a mesoscopic scale over which we average the individual collisions, define our local distribution functions and describe the impact of the collisions on the averaged fluid. Finally, the macroscopic scale on which particles free stream on general relativistic geodesics.

We begin with the description of collisions in the local frame of our observer. The full Hamiltonian H can be separated into a free part H_0 and an interaction part H_I . We employ the Heisenberg picture in which the states are time-independent. The time evolution of our distribution function is given by (omitting to specify the momentum dependence of f_{rs} and N_{rs} for simplicity)

$$\underline{\delta}(0) \frac{d}{dt} f_{rs} = \langle \Psi | \frac{dN_{rs}}{dt} | \Psi \rangle = i \langle \Psi | [H, N_{rs}] | \Psi \rangle = i \langle \Psi | [H_I, N_{rs}] | \Psi \rangle, \quad (98)$$

where in the last identity we have used that helicity is conserved in the absence of collisions since $[H_0, N_{rs}] = 0$. We find a differential equation for the operator N_{rs} and are able to write an approximate solution as closed integration if we restrict ourselves to a given order in the interaction Hamiltonian. We define the operator $N_{rs}^{(0)}$ characterising the ingoing state prior to the collision. To first order in the interaction we obtain

$$N_{rs}(t) = N_{rs}^{(0)} + i \int_0^t dt' [H_I(t'), N_{rs}^{(0)}]. \quad (99)$$

The interpretation of this equation is that at the time $t = 0$ the system is starting to interact, but as the background does not yet contain any correlations between the interacting species we can still evaluate the collisions using the zeroth order number operator. This first order solution describes forward scattering and we need to go to second order to find the first non-forward interactions.

We insert the first order solution (99) into Eq. (98) and find to second order

$$\frac{dN_{rs}}{dt} = i [H_I(t), N_{rs}(t)] \approx i [H_I(t), N_{rs}^{(0)}] - \int_0^t dt' [H_I(t), [H_I(t'), N_{rs}^{(0)}]]. \quad (100)$$

The second order contribution describes an interaction which is active between the time t' and t . We identify this timescale with our microscopic timescale $t_{\text{mic}} = t - t'$, quantifying the timescale of individual particle interactions. The averaged fluid however does not change significantly on this timescale and evolves on the much larger mesoscopic time-scale $t_{\text{mes}} \gg t_{\text{mic}}$. Since we compute the derivative of the number operator N with respect to the time t we may identify the mesoscopic time $t_{\text{mes}} = t$. Expressed in these parameters we obtain

$$\frac{dN_{rs}}{dt_{\text{mes}}} = i [H_I(t_{\text{mes}}), N_{rs}^{(0)}] - \int_0^{t_{\text{mes}}} dt_{\text{mic}} [H_I(t_{\text{mes}}), [H_I(t_{\text{mes}} - t_{\text{mic}}), N_{rs}^{(0)}]] \quad (101)$$

Ideally we would like to evaluate this equation at the initial time and set $t_{\text{mes}} = 0$ to compute the change of our initial states under the considered interactions. This choice however is mathematically inconsistent as we are mixing mesoscopic and microscopic timescales. Instead we average the resulting time-derivative, considering the time-reversal symmetry, over a box that is centered on the initial time and has a length of 2ϵ which is chosen to be small compared to the scale of macroscopic evolution.

$$\left. \frac{dN_{rs}(0)}{dt} \right|_{\text{classical}} \equiv i [H_I(0), N_{rs}^{(0)}] - \int_{-\epsilon}^{\epsilon} \frac{dt_{\text{mes}}}{2\epsilon} \text{sgn}(t_{\text{mes}}) \int_0^{t_{\text{mes}}} dt_{\text{mic}} [H_I(t_{\text{mes}}), [H_I(t_{\text{mes}} - t_{\text{mic}}), N_{rs}^{(0)}]]. \quad (102)$$

We may split this integration into three regions. First, the central region $\epsilon \approx t_{\text{Compton}}$, where t_{Compton} the typical Compton time scale of particles, is highly non-trivial, but this region is negligible compared to our entire integration volume. In the remaining positive and negative regions the integrand is constant in time. The reason is that the integral over the microscopic time already has sufficient support and is converged. The remaining time-dependence based on the mesoscopic time t_{mes} is not relevant as we have chosen the box small compared to the mesoscopic evolution and we may set $t_{\text{mes}} = 0$ yielding

$$\left. \frac{dN_{rs}(0)}{dt} \right|_{\text{classical}} \equiv i [H_I(0), N_{rs}^{(0)}] - \frac{1}{2} \int_{-\epsilon}^{\epsilon} dt_{\text{mic}} [H_I(0), [H_I(t_{\text{mic}}), N_{rs}^{(0)}]]. \quad (103)$$

Finally we may extend the integration limit ϵ to infinity compared to the microscopic evolution using a separation of scales.

We note that the interaction Hamiltonian appearing in this equation may always be evaluated based on the non-interacting field value as we only utilise times which are small compared to the mesoscopic time. Our expression is equivalent to those used in Refs. ([31–33]).

We finally deduce from Eq. (98) that the classical evolution of the distribution function is given by

$$\underline{\delta}(0) \frac{df_{rs}(t)}{dt} = i \langle \Psi(t) | [H_I(t), N_{rs}^{(0)}] | \Psi(t) \rangle - \frac{1}{2} \langle \Psi(t) | \int_{-\infty}^{\infty} dt_{\text{mic}} [H_I(t), [H_I(t + t_{\text{mic}}), N_{rs}^{(0)}]] | \Psi(t) \rangle, \quad (104)$$

where the first term on the rhs is the forward scattering term. It is responsible for refractive effects or flavor oscillations in matter (see Refs. [34, 35] for neutrino oscillations in cosmology) such as the MSW effect [31, 36–38]). The second term is the collision term and we define

$$\underline{\delta}(0) C[f_{rs}(t)] \equiv -\frac{1}{2} \langle \Psi(t) | \int_{-\infty}^{\infty} dt_{\text{mic}} [H_I(t), [H_I(t + t_{\text{mic}}), N_{rs}^{(0)}]] | \Psi(t) \rangle \quad (105)$$

such that the Boltzmann equation (104) (when neglecting forward scattering and restoring the notation of the momentum dependence) is written

$$\underline{\delta}(0) \frac{df_{ss'}(t, p)}{dt} = \underline{\delta}(0) C[f_{ss'}(t, p)]. \quad (106)$$

A spinor space operator associated with this collision term is obtained by contraction with $u_{s'}(p)\bar{u}_s(p)$ (or $v_s(p)\bar{v}_{s'}(p)$ for antiparticles) as in Eq. (38), and we define

$$C[\mathbf{F}(t, p)] \equiv \begin{cases} \sum_{s s'} C[f_{ss'}(t, p)] u_{s'}(p) \bar{u}_s(p), & \text{particles} \\ \sum_{s s'} C[f_{ss'}(t, p)] v_s(p) \bar{v}_{s'}(p), & \text{antiparticles.} \end{cases} \quad (107)$$

The covariant parts of this spinor space collision operator, $I_C(p)$ and $\mathcal{Q}_C^\mu(p)$ are obtained exactly like in Eq. (48). In the massless case the covariant parts are $I_C(p)$, $V_C(p)$ and $Q_C^\mu(p)$ and are obtained as in Eq. (51).

The classical Boltzmann equation is obtained when considering that this derivation, which has been made for a homogeneous system (see § II A), is in fact valid locally. That is in the derivation we assumed that the distribution function depends on time and momentum only $f_{rs}(t, p)$, but we now assume that it also depends on the position and employ $f_{rs}(t, \mathbf{x}, p)$. This amounts to considering that there is a mesoscopic length scale under which the system can be considered as homogeneous such that the volume integral in the Hamiltonian $H_I = \int d^3\mathbf{x} \mathcal{H}_I$ can be extended to infinity in the computation of the local collision term $C[f_{ss'}(t, \mathbf{x}, p)]$. Expressed in terms of spinor valued operators the classical Boltzmann equation reads

$$\frac{d\mathbf{F}(t, \mathbf{x}, p)}{dt} = C[\mathbf{F}(t, \mathbf{x}, p)]. \quad (108)$$

B. General relativity and the classical Boltzmann equation in curved space-time

We connect the collision term derived in the mesoscopic Minkowski space-time to the macroscopic relativistic evolution. We assume that collisions are well described in special relativity and general relativistic corrections may be neglected. Under our assumptions, the covariant parts of the local spinor space collision operator (107), do also depend on the position of space [e.g. we should write $I_C(t, \mathbf{x}, p)$]. Using $dt/d\lambda = p^0 = E$, the derivative $E dI/dt$ is identified with $DI/D\lambda$ of the parallel transport equation (86), but now the effect of collisions is taken into account by the intensity part of Eq. (108), $dI/dt = I_C$. This means that we connect the time in the local inertial frame, in which we compute the collisions, with the time along our general relativistic geodesic.

Similarly, expressing the polarisation part of the collision term in general coordinates as in Eq. (87), that is with

$$\mathcal{Q}_C^\alpha = \mathcal{Q}_C^\mu [e_\mu]^\alpha, \quad (109)$$

we identify $E d\mathcal{Q}^\alpha/dt$ with $D\mathcal{Q}^\alpha/D\lambda$ of the parallel transport equation (88) and we find

$$\frac{DI}{D\lambda} = E I_C, \quad \frac{D\mathcal{Q}^\alpha}{D\lambda} = E \mathcal{Q}_C^\alpha. \quad (110)$$

This is the general relativistic Boltzmann equation for fermions, needed to compute the effect of both free streaming and collisions on the distribution function of fermions. When computing the collision term for particular examples in § VI, it is convenient to present the results with a prefactor $1/E$ so that the rhs of Eq. (110) can be readily obtained.

When expressed in functions of the general time coordinate τ , we just use $d\tau/d\lambda = P^0$ to deduce that the rhs of Eqs. (89) and (90) are respectively $E/P^0 I_C$ and $E/P^0 \mathcal{Q}_C^\alpha$ when collisions are to be considered.

C. Flavor description and flavor oscillations

We will now consider the case of flavoured particles, following Ref. [31]. We add flavour indices ($i, j = 1, \dots, n_{\text{flavour}}$) to our creation and annihilation operators a_s^i and the distribution function $f_{r_i s_j}$ is obtained from $a_r^{\dagger i} a_s^j$ following the same procedure as in §II A. For simplicity of notation, we omit the spin indices r, s in this section. These flavour-states are chosen to be diagonal in the interactions, but the corresponding mass matrix may not be diagonal and thus the flavour states are no longer conserved in free propagation.

In our approach the oscillations are represented by forward scatterings. The free Hamiltonian takes the form

$$H_0 = \int [dp] \sum_{ij} \sum_s a_s^{i\dagger}(p) \Omega_{ij} a_s^j(p) + b_s^{i\dagger}(p) \Omega_{ij} b_s^j(p), \quad (111)$$

with $\Omega_{ij} = \sqrt{\mathbf{p}^2 + M_{ij}^2}$ a symmetric matrix and M_{ij} the neutrino mass matrix. In the absence of interactions we find⁴

$$\frac{dN_{ij}^{(0)}(t)}{dt} = i[H_0(t), N_{ij}^{(0)}]. \quad (112)$$

where the commutator between the free Hamiltonian and the number operator does not vanish. We explicitly obtain

$$\frac{df_{ij}}{dt} = i \sum_k (\Omega_{ik} f_{kj} - f_{ik} \Omega_{kj}). \quad (113)$$

When considering both interactions and flavour oscillations, we assume a separation of scales between the flavour oscillations, relevant on the mesoscopic scale, and the collisions on the microscopic scale. This means that we allow for flavour oscillations in-between two individual interactions, but not during one single collision. This assumption allows us to write up to second order in collisions

$$\begin{aligned} \underline{\delta}(0) \frac{df_{ij}(t)}{dt} &= i \langle \Psi(t) | [H_0(t), N_{ij}] | \Psi(t) \rangle + i \langle \Psi(t) | [H_I(t), N_{ij}^{(0)}] | \Psi(t) \rangle \\ &\quad - \frac{1}{2} \langle \Psi(t) | \int_{-\infty}^{\infty} dt_{\text{mic}} [H_I(t), [H_I(t + t_{\text{mic}}), N_{ij}^{(0)}]] | \Psi(t) \rangle. \end{aligned} \quad (114)$$

Defining the collision term as in Eq. (105) and neglecting forward scattering from the interaction Hamiltonian, we obtain the classical Boltzmann equation

$$\frac{df_{ij}(t, p)}{dt} = i \sum_k (\Omega_{ik}(p) f_{kj}(t, p) - f_{ik}(t, p) \Omega_{kj}(p)) + C[f_{ij}(t, p)]. \quad (115)$$

Note that only the spatial momentum is conserved (and not energy of particles, because they change from one mass shell to another one) by the flavor changing term but we must not forget that when we consider a distribution function f_{ij} we must extract its energy with $\sum_{ij} \Omega_{ij} f_{ij}$ and it is easily seen that this is conserved by the flavor changing term. To promote the Boltzmann equation to general relativity we follow the recipe presented in the previous sections. We construct the observer-independent quantities I_{ij} and Q_{ij} from the distribution function f_{ij} and connect them to general relativistic manifold by multiplying them with the tetrad. The flavour oscillations can be treated in the same way as the collision term. They are well described on the mesoscopic scale, independent from the geometrical corrections of general relativity relevant on the macroscopic scale. As particles are propagating on the free geodesics of general relativity, the flavour of the particles is mixed. We therefore connect the time measured by the local observer describing the flavour oscillations to our relativistic time coordinate and, similarly to section VB we find that the factor E/P_0 must be added in front of the flavour oscillations term [the first term on the rhs of Eq. (115)] before incorporating it to the rhs of the Boltzmann equations (89) and (90).

⁴ Note that in Eq. (2.5) of Ref. [31] there is an extra minus sign for particles compared to antiparticles because the definition of the distribution function is the transposed of our definition f_{rs} .

VI. COLLISIONS MEDIATED BY WEAK INTERACTIONS

In order to illustrate how the previous formalism should be implemented in practice, we will focus on weak-interactions, and more precisely on their low-energy limit. Since this has the advantage of involving only fermions it is very well suited for the formalism introduced in this article.

A. General form of weak-interactions

All weak interaction take the form of current-current interactions [39] at low energy (low compared to the W^\pm and Z masses), that is they are given by

$$\mathcal{H}_I = -\mathcal{L}_I = -\frac{G_F}{\sqrt{2}} [J_\mu^{\text{NC}} J_{\text{NC}}^\mu + J_\mu^{\text{CC}\dagger} J_{\text{CC}}^\mu], \quad (116)$$

where $G_F \simeq 1.17 \times 10^{-5} \text{ GeV}^{-2}$ is the Fermi constant of weak interactions.

1. Neutral currents

Neutral currents describe the exchange of Z bosons and as these are not charged they mediate elastic scatterings that do not alter the involved types of particles and only transfer momentum, spin and energy.

The neutral current is simply the sum of the neutral currents of all particles undergoing weak interactions

$$J_{\text{NC}}^\mu = J_{ee}^\mu + J_{\nu\nu}^\mu + \dots \quad (117)$$

For neutrinos, the neutral current couples only the left chiralities and, noting ν the neutrino quantum field, it is simply

$$J_{\nu\nu}^\mu \equiv e_-^\nu \bar{\nu} \gamma^\mu (\mathbb{1} - \gamma_5) \nu, \quad e_-^\nu \equiv \frac{1}{2}. \quad (118)$$

with similar expressions for other flavors. However for electrons and (similarly pions and taus) the neutral currents must be further decomposed into left and right chiral interactions as

$$J_{ee}^\mu = \epsilon_-^e J_{ee}^{-\mu} + \epsilon_+^e J_{ee}^{+\mu}, \quad J_{ee}^{-\mu} \equiv \bar{e} \gamma^\mu (\mathbb{1} - \gamma_5) e, \quad J_{ee}^{+\mu} \equiv \bar{e} \gamma^\mu (\mathbb{1} + \gamma_5) e, \quad (119)$$

where we noted e the electronic quantum field. The chiral coupling constants are for electrons

$$\epsilon_-^e \equiv -\frac{1}{2} + \sin^2 \theta_W, \quad \epsilon_+^e \equiv \sin^2 \theta_W, \quad (120)$$

with θ_W the Weinberg angle ($\sin^2 \theta_W \simeq 0.23$).

2. Charged currents

Opposed to the neutral currents, the charged currents describe the exchange of charged W -bosons and therefore are inelastic. The structure of the charged currents is more complex since they couple eigenmass states of different flavors, thanks to the Cabbibo-Kobayashi-Maskawa (CKM) matrix for quarks or the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix for massive neutrinos. We ignore these complications for the examples that we shall consider and consider only the effective charged currents for the neutron/proton pair which is involved in beta decays and related processes, and the charged currents of the first two lepton flavors, that is of the electron/neutrino and muon/muon neutrino pairs.

$$J_{CC}^\sigma = \cos \theta_C J_{pn}^\sigma + J_{e\nu}^\sigma + J_{\mu\nu\mu}^\sigma, \quad (121a)$$

$$(121b)$$

where $\cos \theta_C = 0.98$ is a Cabbibo-Kobayashi-Maskawa (CKM) angle.

The charged currents for electron/neutrino and muon/muon neutrino pairs are coupling only the left chiralities

$$J_{e\nu}^\sigma \equiv \bar{\nu}\gamma^\sigma(\mathbb{1} - \gamma^5)\mathbf{e}, \quad J_{\mu\nu\mu}^\sigma \equiv \bar{\nu}_\mu\gamma^\sigma(\mathbb{1} - \gamma^5)\boldsymbol{\mu}. \quad (122)$$

However, due to internal QCD effects, the coupling in the proton/pair is not purely left chiral. The deviation from left chirality of the coupling is parameterized by the g_A parameter whose measured value is approximately 1.25 [39] and the corresponding charged current reads

$$J_{pn}^\mu \equiv \bar{\mathbf{p}}\gamma^\mu(\mathbb{1} - g_A\gamma^5)\mathbf{n}. \quad (123)$$

When considering the cumulative effect of neutral currents and charged currents, we can use the Fierz identities which for anticommuting fields give [31, 40]

$$J_{e\nu}^{\dagger\mu}(J_{e\nu})_\mu = J_{ee}^{-\mu}(J_{\nu\nu}^-)_\mu. \quad (124)$$

This means that the effect of the charged current can be replaced by an equivalent neutral current. Instead of computing the collision term for charged currents, we may alternatively modify the neutral currents chiral coupling factors (120), yielding

$$\epsilon_-^e \rightarrow \epsilon_-^e + 1. \quad (125)$$

B. General current/current interaction

For any reaction involving weak-interactions, a visual inspection of the interaction Hamiltonian is sufficient to deduce which currents are involved in the process. In all cases this amounts to considering the current-current coupling between four species a, b, c, d given by

$$\mathcal{H}_I = -\mathcal{L}_I = -g(J_\mu^{ac}J_{bd}^\mu + cc), \quad (126)$$

where, depending on the interaction, the same species may be represented by multiple indices. The chiral contributions of these currents are parameterized by ϵ_\pm^{ac} and ϵ_\pm^{bd} as

$$J_{ac}^\mu = \bar{\psi}_c\chi_{(ac)}^\mu\psi_a, \quad J_{bd}^\mu = \bar{\psi}_d\chi_{(bd)}^\mu\psi_b, \quad (127)$$

with the notation

$$\chi_{(ac)}^\mu \equiv 2 \sum_{s=\pm 1} \epsilon_s^{ac}\gamma^\mu P_s = \epsilon_+^{ac}\gamma^\mu(\mathbb{1} + \gamma^5) + \epsilon_-^{ac}\gamma^\mu(\mathbb{1} - \gamma^5) \quad (128a)$$

$$\chi_{(bd)}^\mu \equiv 2 \sum_{s=\pm 1} \epsilon_s^{bd}\gamma^\mu P_s = \epsilon_+^{bd}\gamma^\mu(\mathbb{1} + \gamma^5) + \epsilon_-^{bd}\gamma^\mu(\mathbb{1} - \gamma^5). \quad (128b)$$

We now study the structure of the collision term due the general interaction Hamiltonian (126) with the currents (127), and we apply it further to specific cases, which all correspond to a choice of the species a, b, c, d together with the couplings ϵ_\pm^{ac} and ϵ_\pm^{bd} and the coupling constant g .

Let us investigate the total collision term for the particle species a due to this interactions, which is represented by the sum of collision terms. First there is the decay of particle a corresponding to the reaction

$$a \rightarrow \bar{b} + c + d \quad (129)$$

where we recall that \bar{b} is the antiparticle species related to the particle species b . We may typically neglect the three body reactions that would revert this decay, as they are only relevant in very high density environments. In addition to the decay, we have to consider the two-body reactions

$$a + b \leftrightarrow c + d \quad (130a)$$

$$a + \bar{c} \leftrightarrow \bar{b} + d \quad (130b)$$

$$a + \bar{d} \leftrightarrow c + \bar{b}. \quad (130c)$$

However all the related collision terms can be deduced from the one of $a + b \leftrightarrow c + d$ through crossing symmetry and we discuss this procedure in the next section. Here we focus only on this specific reaction.

C. General method for the generic process $a + b \leftrightarrow c + d$

Let us introduce some compact notation with the multi-indices

$$\alpha \equiv (s_\alpha, p_\alpha) \quad \alpha' \equiv (s'_\alpha, p'_\alpha). \quad (131)$$

These multi-indices contain all information characterising one single particle (its momentum and helicity). We will typically label ingoing states as unprimed and outgoing states with primed indices. For species a we employ the multi-index α and similarly for species b (resp. c and d) we use the multi-indices β (resp. γ and δ). The plane wave solutions are written in a compact form in this notation. For instance for the species a we write $u_\alpha \equiv u_{s_\alpha}(p_\alpha)$ and $v_\alpha \equiv v_{s_\alpha}(p_\alpha)$. Furthermore this allows to write a compact relativistic Dirac delta function which acts both on helicities and momenta as

$$\underline{\delta}_{\alpha\alpha'} \equiv \delta_{s_\alpha s'_\alpha}^K \underline{\delta}(p_\alpha - p'_\alpha). \quad (132)$$

We denote the number operator associated with species a as

$$A_{\alpha\alpha'} \equiv N_{s_\alpha s'_\alpha}(p_\alpha, p'_\alpha) = a_{s_\alpha}^\dagger(p_\alpha) a_{s'_\alpha}(p'_\alpha). \quad (133)$$

We also define the Pauli blocking operator

$$\widehat{A}_{\alpha\alpha'} \equiv \underline{\delta}_{\alpha\alpha'} - A_{\alpha\alpha'}. \quad (134)$$

The expectation value of these operators is denoted as

$$\langle A_{\alpha\alpha'} \rangle = \underline{\delta}(p_\alpha - p'_\alpha) \mathcal{A}_{\alpha\alpha'}(p_\alpha), \quad \langle \widehat{A}_{\alpha\alpha'} \rangle = \underline{\delta}(p_\alpha - p'_\alpha) \widehat{\mathcal{A}}_{\alpha\alpha'}(p_\alpha) \quad (135)$$

where we introduce the short-hand notation $\mathcal{A}_{\alpha\alpha'}(p_\alpha) = \mathcal{A}_{s_\alpha s'_\alpha}(p_\alpha)$. We recall that this quantity is exactly the one-particle distribution function associated with species a . Note that for the Pauli blocking factor, $\widehat{\mathcal{A}}_{\alpha\alpha'}(p_\alpha)$ is a shorthand notation for $\delta_{s_\alpha s'_\alpha}^K - \mathcal{A}_{s_\alpha s'_\alpha}(p_\alpha)$. We associate to the one-particle distribution function (resp. the Pauli blocking function) a spinor valued operator following the procedure (38) that we note A_a^b (resp. \widehat{A}_a^b) in component notation or simply \mathbf{A} (resp. $\widehat{\mathbf{A}}$) in operator notation. Having defined for species a the number operator $A_{\alpha\alpha'}$, the distribution function $\mathcal{A}_{\alpha\alpha'}$ and the spinor-valued (observer-independent) operator \mathbf{A} , identically for species b (resp. c , d) we use $B_{\beta\beta'}$, $\mathcal{B}_{\beta\beta'}$ and \mathbf{B} (resp. $C_{\gamma\gamma'}$, $\mathcal{C}_{\gamma\gamma'}$ and \mathbf{C} , $D_{\delta\delta'}$, $\mathcal{D}_{\delta\delta'}$ and \mathbf{D}), and associated hatted notations for Pauli blocking factors. Furthermore, for the antiparticles species \bar{a} , \bar{b} , \bar{c} , \bar{d} related to the species a , b , c , d , we use barred notation for number operators (e.g. $\bar{A}_{\alpha\alpha'}$), distribution function (e.g. $\bar{\mathcal{A}}_{\alpha\alpha'}$) and spinor valued operators (e.g. $\bar{\mathbf{A}}$), along with their hatted versions for Pauli blocking terms. Finally we define the collision term as in Eq. (106), that is

$$\underline{\delta}(0) C[\mathcal{A}_{ss'}(p)] \equiv -\frac{1}{2} \left\langle \int_{-\infty}^{\infty} dt' [H_I(0), [H_I(t'), \mathcal{A}_{ss'}(p)]] \right\rangle \quad (136)$$

such that the quantum Boltzmann equation (104) for species a is written as (when neglecting forward scattering)

$$\underline{\delta}(0) \frac{d\mathcal{A}_{ss'}(p)}{dt} = \underline{\delta}(0) C[\mathcal{A}_{ss'}(p)]. \quad (137)$$

Following the previous discussion, our goal is to compute the collision term $C[\mathcal{A}_{ss'}(p)]$ corresponding to the reaction $a + b \leftrightarrow c + d$, when considering an interaction Hamiltonian of the form (126). This interaction Hamiltonian contains the term

$$H_I^{a+b \leftrightarrow c+d} \equiv \int [dp_\alpha][dp_\beta][dp_\gamma][dp_\delta] (2\pi)^3 \delta(\mathbf{p}_\alpha + \mathbf{p}_\beta - \mathbf{p}_\gamma - \mathbf{p}_\delta) e^{-i(p_\alpha^0 + p_\beta^0 - p_\gamma^0 - p_\delta^0)t} \mathcal{M}^{a+b \leftrightarrow c+d} \quad (138)$$

which is describing the reaction $a + b \leftrightarrow c + d$, and where we used the scattering operator for this reaction

$$\mathcal{M}^{a+b \leftrightarrow c+d} \equiv \sum_{\text{spins}} \left(d_\delta^\dagger c_\gamma^\dagger b_\beta a_\alpha M_{\alpha\beta \rightarrow \gamma\delta} + b_\beta^\dagger a_\alpha^\dagger d_\delta c_\gamma M_{\gamma\delta \rightarrow \alpha\beta} \right). \quad (139)$$

The M matrices are

$$M_{\alpha\beta \rightarrow \gamma\delta} \equiv M[(s_\alpha, p_\alpha)(s_\beta, p_\beta) \rightarrow (s_\gamma, p_\gamma)(s_\delta, p_\delta)] \equiv g[\bar{u}_\gamma \chi_{(ac)}^\mu u_\alpha][\bar{u}_\delta \chi_\mu^{(bd)} u_\beta] \quad (140a)$$

$$M_{\gamma\delta \rightarrow \alpha\beta} = M_{\alpha\beta \rightarrow \gamma\delta}^* = g[\bar{u}_\alpha \chi_{(ac)}^\mu u_\gamma][\bar{u}_\beta \chi_\mu^{(bd)} u_\delta]. \quad (140b)$$

To compute the collision term we first need to compute the operator $[\mathcal{M}, [\mathcal{M}, A_{ss'}]]$. Using the commutation rules given in Appendix B, we get

$$\begin{aligned} [\mathcal{M}, [\mathcal{M}, A_{ss'}]] &\equiv M_{\gamma'\delta'\rightarrow\alpha\beta} M_{\alpha'\beta'\rightarrow\gamma\delta} \left\{ [d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} a_{\alpha}^{\dagger}, d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{s'}] [a_{\alpha'}, a_s^{\dagger} a_{s'}] + [d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{\alpha'}, d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} [a_{\alpha}^{\dagger}, a_s^{\dagger} a_{s'}]] \right\} \\ &= M_{\gamma'\delta'\rightarrow\alpha\beta} M_{\alpha'\beta'\rightarrow\gamma\delta} \left\{ [d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} a_{\alpha}^{\dagger}, d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{s'}] \underline{\delta}_{s\alpha'} - [d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{\alpha'}, d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} a_s^{\dagger}] \underline{\delta}_{s'\alpha} \right\} \\ &= M_{\gamma'\delta'\rightarrow\alpha\beta} M_{\alpha'\beta'\rightarrow\gamma\delta} \left\{ [d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} a_{\alpha}^{\dagger}, d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{s'}] \underline{\delta}_{s\alpha'} + [d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} a_s^{\dagger}, d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{\alpha'}] \underline{\delta}_{s'\alpha} \right\}. \end{aligned} \quad (141)$$

The commutators of this last expression are expressible simply in terms of the number operators of the species. For instance using the commutation rules of Appendix B we get

$$[d_{\delta'} c_{\gamma'} b_{\beta}^{\dagger} a_{\alpha}^{\dagger}, d_{\delta}^{\dagger} c_{\gamma}^{\dagger} b_{\beta'} a_{s'}] = A_{\alpha s'} B_{\beta\beta'} \widehat{C}_{\gamma\gamma'} \widehat{D}_{\delta\delta'} - \widehat{A}_{\alpha s'} \widehat{B}_{\beta\beta'} C_{\gamma\gamma'} D_{\delta\delta'}. \quad (142)$$

Hence we obtain

$$\begin{aligned} [\mathcal{M}, [\mathcal{M}, A_{ss'}]] &= M_{\alpha\beta\rightarrow\gamma'\delta'}^* M_{\alpha'\beta'\rightarrow\gamma\delta} \\ &\left\{ \underline{\delta}_{s\alpha'} \left[B_{\beta\beta'} A_{\alpha s'} \widehat{D}_{\delta\delta'} \widehat{C}_{\gamma\gamma'} - D_{\delta\delta'} C_{\gamma\gamma'} \widehat{B}_{\beta\beta'} \widehat{A}_{\alpha s'} \right] + \underline{\delta}_{\alpha s'} \left[B_{\beta\beta'} A_{s\alpha'} \widehat{D}_{\delta\delta'} \widehat{C}_{\gamma\gamma'} - D_{\delta\delta'} C_{\gamma\gamma'} \widehat{B}_{\beta\beta'} \widehat{A}_{s\alpha'} \right] \right\}. \end{aligned} \quad (143)$$

We now employ this result in Eqs. (138) and (136). We integrate a total of five momentum integrals (each one being itself three-dimensional in momentum space) using the Dirac distributions. Of these, four Dirac functions are contained in the expectation values of the number operators associated to the four species, and there is an extra Dirac function $\underline{\delta}_{s\alpha'}$ from the collision term ensuring local energy and momentum conservation. Eventually, taking the expectation in the quantum state, we get

$$\begin{aligned} \underline{\delta}(0) C[\mathcal{A}_{ss'}(p)] &= \mathcal{V} \mathcal{K} M^* [(s_{\alpha}, p) (s_{\beta}, p_{\beta}) \rightarrow (s'_{\gamma}, p_{\gamma}) (s'_{\delta}, p_{\delta})] M [(s'_{\alpha}, p) (s'_{\beta}, p_{\beta}) \rightarrow (s_{\gamma}, p_{\gamma}) (s_{\delta}, p_{\delta})] \\ &\left\{ \delta_{s\alpha'}^{\mathcal{K}} \left[-\mathcal{B}_{\beta\beta'}(p_{\beta}) \mathcal{A}_{\alpha s'}(p) \widehat{D}_{\delta\delta'}(p_{\delta}) \widehat{C}_{\gamma\gamma'}(p_{\gamma}) + \mathcal{D}_{\delta\delta'}(p_{\delta}) \mathcal{C}_{\gamma\gamma'}(p_{\gamma}) \widehat{B}_{\beta\beta'}(p_{\beta}) \widehat{A}_{\alpha s'}(p) \right] \right. \\ &\left. + \delta_{\alpha s'}^{\mathcal{K}} \left[-\mathcal{B}_{\beta\beta'}(p_{\beta}) \mathcal{A}_{s\alpha'}(p) \widehat{D}_{\delta\delta'}(p_{\delta}) \widehat{C}_{\gamma\gamma'}(p_{\gamma}) + \mathcal{D}_{\delta\delta'}(p_{\delta}) \mathcal{C}_{\gamma\gamma'}(p_{\gamma}) \widehat{B}_{\beta\beta'}(p_{\beta}) \widehat{A}_{s\alpha'}(p) \right] \right\} \end{aligned} \quad (144)$$

with the integration on momenta

$$\mathcal{K} = \frac{1}{2} \int [dp_{\delta}] [dp_{\gamma}] [dp_{\beta}] (2\pi)^4 \delta^{(4)}(p_{\delta} + p_{\gamma} - p_{\beta} - p). \quad (145)$$

We note that:

- The collision term is made of two types of terms. The first terms on the second and the third line of Eq. (144) correspond to *scattering out* processes, that is collisions which due to the minus sign deplete the distribution function associated with species a and they correspond to $a + b \rightarrow c + d$. The second term on the second and third line correspond conversely to *scattering in* processes, which increase the distribution function of species a , and they represent interactions $c + d \rightarrow a + b$.
- For scattering out processes, the collision term is proportional to the distribution function of the initial states (species a and b), but also to the Pauli blocking function of the final states (species c and d), and the reverse is true for the scattering in processes.
- The distribution functions are Hermitian, that is $\mathcal{A}_{s s'}^*(p) = \mathcal{A}_{s' s}(p)$ as in Eq. (27). Let us now consider $C[\mathcal{A}_{ss'}(p)]^*$. Given the Hermiticity of the distribution functions and thus of the Pauli blocking functions, then with a simple renaming of all primed indices as unprimed indices (and also of unprimed indices as primed indices), it is straightforward to show that this is equal to $C[\mathcal{A}_{s' s}(p)]$, hence the collision term is also Hermitian as expected.
- In the previous computation when checking the Hermiticity, the second and third line of Eq. (144) are interchanged. Terms of the second line are proportional to $\delta_{s\alpha'}^{\mathcal{K}}$ and correspond physically to the scattering of the helicity index s' , and conversely in the third line the terms are proportional to $\delta_{\alpha s'}^{\mathcal{K}}$ and it corresponds to the scattering of the helicity index s . Hence we see that the collision term possesses four terms corresponding to the *in/out* contributions and the s/s' contributions.

- Finally even though we computed the collision term for a homogenous system in a Minkowski space-time, the total volume which appears as $\delta^{(3)}(0)$ drops out from both the left and the right hand side of Eq. (144) and thus of Eq. (137). Hence, as argued before Eq. (108), we can consider that this collision term is valid locally, allowing us to consider in a classical macroscopic description that all distribution functions should be considered with a dependence on the point of space-time. We started a computation with total number of particles in a quantum system, but we end up using it with *number densities* of particles, considering that the collisions are point-like.

The procedure to follow is now transparent. The helicity indices of the distribution functions (or the related Pauli blocking functions) are contracted with the plane waves solutions contained in the M matrices. From Eqs. (38) this is exactly what is needed to build the spinor space operators related to each species. Since only the indices s and s' remain uncontracted in Eq. (144), we contract them with $u_{s'}(p)\bar{u}_s(p)$ (or $v_s(p)\bar{v}_{s'}(p)$ for antiparticles) so as to form a spinor space collision operator $C[\mathbf{A}(p)]$ as specified in the definition (107). Note that the contraction of $\delta_{\alpha s'}^K$ with $u_{s'}(p)\bar{u}_s(p)$ or $v_s(p)\bar{v}_{s'}(p)$ gives simply $-\not{p} + M$ as can be seen from Eqs. (43).

We finally obtain

$$C[\mathbf{A}(p)] = -\frac{g^2}{2E_a} \mathcal{K} \left\{ \text{Tr}[\mathbf{B} \cdot \chi_\mu^{(bd)} \cdot \hat{\mathbf{D}} \cdot \chi_\nu^{(bd)}] \times \left[\mathbf{A} \cdot \chi_{(ac)}^\mu \cdot \hat{\mathbf{C}} \cdot \chi_{(ac)}^\nu \cdot (-\not{p} + M) + (-\not{p} + M) \cdot \chi_{(ac)}^\mu \cdot \hat{\mathbf{C}} \cdot \chi_{(ac)}^\nu \cdot \mathbf{A} \right] - (\mathbf{A} \leftrightarrow \hat{\mathbf{A}}, \mathbf{B} \leftrightarrow \hat{\mathbf{B}}, \hat{\mathbf{C}} \leftrightarrow \mathbf{C}, \hat{\mathbf{D}} \leftrightarrow \mathbf{D}) \right\} \quad (146)$$

where the momentum dependence $\mathbf{A}(p)$ and $\mathbf{B}(p_\beta)$, $\mathbf{C}(p_\gamma)$, $\mathbf{D}(p_\delta)$ (and similarly for Pauli blocking operators) are omitted for a more compact notation. The structure of the collision term is again manifest. The first line corresponds to scattering out processes. It is made of two terms which thanks to the property (53) of all operators appearing ($\{\mathbf{A}, \mathbf{B}, \hat{\mathbf{C}}, \hat{\mathbf{D}}, \chi_{(ac)}^\mu, \chi_{(bd)}^\mu\}$), ensure that the property (53) is also satisfied for the collision operator. As for the second line, it corresponds to the scattering in processes, and differs only by an overall sign and the exchange of the distribution and Pauli blocking functions.

This collision term $C[\mathbf{A}]$, being itself an operator in spinor space, can be decomposed into its covariant parts $I_{C[\mathbf{A}]}$ and $\mathcal{Q}_{C[\mathbf{A}]}$ as in the decomposition (48). These components can be found by multiplying by the appropriate $X \in \mathcal{O}$ and taking the trace, that is using the extractions (55) or (57) in the massless case. Since all operators involved in the collision term are made of γ^μ or γ^5 matrices, the problem is reduced to taking traces of products of these operators. The corresponding expressions are gathered in Appendix C. This procedure is conceptually simple and standard, but it can become rather involved when there is a large number of operators in the products. Indeed, letting aside the γ^5 operators, the largest products involve two γ^μ matrices for each distribution function or Pauli blocking function from the $\tilde{\Sigma}_{\mu\nu}$. Then considering the γ^μ of the operators (128), and the one contained in $-\not{p} + M$, we see that when extracting the components on γ^μ (resp. $\Sigma_{\mu\nu}$) of the collision term, we end up with a total of eight (resp. nine) γ^μ operators. However, this systematic computation can be handled by a computer algebra package such as *xAct* [41] and this is particularly powerful since it also takes care of all simplifications involving space-time indices.

In particular, when using Eqs. (36) to extract the intensity part of the collision term (146), we find using the decomposition (50) for $\mathbf{A}(p)$ and the property $(\not{p} + M)(\not{p} + M) = 2M(\not{p} + M)$ (deduced from $\not{p}\not{p} = -m^2$) its general expression

$$\frac{1}{2} I_{C[\mathbf{A}]}(p) = -\frac{g^2}{2E_a} \mathcal{K} \left\{ \text{Tr}[\mathbf{B} \cdot \chi_\mu^{bd} \cdot \hat{\mathbf{D}} \cdot \chi_\nu^{bd}] \text{Tr} \left[\mathbf{A} \cdot \chi_{ac}^\mu \cdot \hat{\mathbf{C}} \cdot \chi_{ac}^\nu \right] - (\mathbf{A} \leftrightarrow \hat{\mathbf{A}}, \mathbf{B} \leftrightarrow \hat{\mathbf{B}}, \hat{\mathbf{C}} \leftrightarrow \mathbf{C}, \hat{\mathbf{D}} \leftrightarrow \mathbf{D}) \right\}. \quad (147)$$

This general expression is also valid in the massless case. To show this we first need that for any two four-vectors A^μ and B^μ , $\{\hat{\mathbf{A}}, \hat{\mathbf{B}}\} = -2A^\mu B_\mu \mathbf{1}$ thanks to Eq. (11), so that $\{\not{p}, \not{k}\} = 2\mathbf{1}$ and $\{\not{p}, \not{Q}\} = 0$. Using the extraction expression (57a) applied on the collision term (146), and commuting the operators with the previous properties to force the appearance of $\not{p}\not{p} = 0$, leads also to Eq. (147).

The collision operator (146) has the general form

$$C[\mathbf{A}(p)] = -[(\not{p} + M) \cdot \mathbf{K} \cdot \mathbf{A}(p) + \mathbf{A}(p) \cdot \mathbf{K} \cdot (\not{p} + M)] + [(\not{p} + M) \cdot \hat{\mathbf{K}} \cdot \hat{\mathbf{A}}(p) + \hat{\mathbf{A}}(p) \cdot \hat{\mathbf{K}} \cdot (\not{p} + M)], \quad (148)$$

where $\mathbf{K} = \mathbf{K}[\mathbf{B}, \hat{\mathbf{C}}, \hat{\mathbf{D}}]$ is an operator [satisfying property (53)] depending on other species distribution functions integrated over momenta, and $\hat{\mathbf{K}} = \mathbf{K}[\hat{\mathbf{B}}, \mathbf{C}, \mathbf{D}]$ is its hatted version. For other types of interactions than the current-current weak interactions that we considered (e.g. considering the effect of the finite mass of the vector boson exchanged), the collision operator for the species a would still have the general form (148) but with a different \mathbf{K} . If the interactions are with bosons, the hatted expressions in \mathbf{K} have to be understood in the sense of stimulated

emission instead of Pauli blocking as discussed in Appendix A. From the general form (148), the intensity part is in general obtained from

$$\frac{1}{2}I_{C[A]}(p) = -\text{Tr}[\mathbf{K} \cdot \mathbf{A}(p)] + \text{Tr}[\widehat{\mathbf{K}} \cdot \widehat{\mathbf{A}}(p)]. \quad (149)$$

D. Explicit form of the collision term for $a + b \leftrightarrow c + d$

1. Notation

Let us introduce some notation which allows to give explicitly the collision term for the species a corresponding to the reaction $a + b \leftrightarrow c + d$, as obtained from the method exposed in the previous section. From the covariant components of the distribution function which are $I(p)$ and $Q^\mu(p)$, we can build scalar, vectors and tensors with which the collision term is better expressed. Let us first introduce the chiral vectors $\mathcal{I}_\mu^\pm(p)$ (see appendix (D 8) for physical interpretation) and the achiral vector $\mathcal{I}_\mu^0(p)$

$$2\mathcal{I}_\mu^\pm \equiv Ip_\mu \pm MQ_\mu = Ip_\mu \pm MQ_\mu \pm MVS_\mu, \quad 2\mathcal{I}_\mu^0 \equiv Ip_\mu, \quad (150)$$

where we remind the notation $M = m$ (resp. $M = -m$) for a particle (resp. for an antiparticle) and where the dependence of all quantities on the momentum p^μ is omitted. Note that the left chiral vector for a particle \mathcal{I}_μ^- is equal to the right chiral vector \mathcal{I}_μ^+ of the corresponding antiparticle and vice versa. We also define the Pauli blocking vectors

$$\widehat{\mathcal{I}}_\mu^\pm \equiv p_\mu - \mathcal{I}_\mu^\pm, \quad \widehat{\mathcal{I}}_\mu^0 \equiv p_\mu - \mathcal{I}_\mu^0, \quad \widehat{Q}^\mu = -Q^\mu. \quad (151)$$

In the massless case, using that $mS^\mu \rightarrow p^\mu$ we must take the definitions

$$2\mathcal{I}_\mu^\pm \equiv (I \pm \lambda V)p_\mu. \quad (152)$$

where we remind that λ is 1 for particles and -1 for antiparticles. Note that the purely linear polarisation Q^μ does not contribute in the definitions of the chiral vectors in the massless case. We then define scalar quantities

$$\mathcal{I} \equiv \frac{I}{2}, \quad \widehat{\mathcal{I}} = 1 - \mathcal{I}, \quad \mathcal{J} \equiv M\mathcal{I} = M\frac{I}{2}, \quad \widehat{\mathcal{J}} \equiv M\widehat{\mathcal{I}} = M\left(1 - \frac{I}{2}\right), \quad (153)$$

and tensor quantities

$$\mathcal{S}_{\mu\nu} \equiv p_{[\mu}Q_{\nu]} \quad \widehat{\mathcal{S}}_{\mu\nu} = -\mathcal{S}_{\mu\nu} \quad (154a)$$

$$\mathcal{S}_{\mu\nu}^\pm = \mathcal{S}_{\mu\nu} \mp \mathcal{J}g_{\mu\nu} = \mathcal{S}_{\mu\nu} \mp \frac{MI}{2}g_{\mu\nu}, \quad (154b)$$

where again the hatted notation refers to Pauli blocking forms. In the massless limit $\mathcal{J} = \widehat{\mathcal{J}} = 0$ and $\mathcal{S}_{\mu\nu}^\pm = \mathcal{S}_{\mu\nu}$. Furthermore $\mathcal{S}_{\mu\nu} = p_{[\mu}Q_{\nu]}$, and it is non-vanishing only if there is linear polarisation.

The collision term is expressed by means of various contractions of these scalars, vectors and tensors associated with the various species involved in the collision. In order to obtain a compact result, we introduce the following contractions (we remind that indices a, b, c, d refer to the species considered):

$$[\mathcal{S}_a \cdot \mathcal{S}_b] \equiv \mathcal{S}_{a\nu}^\mu \mathcal{S}_{b\mu}^\nu, \quad [\mathcal{S}_a \cdot \mathcal{S}_b \cdot \mathcal{S}_c \cdot \mathcal{S}_d] \equiv \mathcal{S}_{a\lambda}^\sigma \mathcal{S}_{b\mu}^\lambda \mathcal{S}_{c\nu}^\mu \mathcal{S}_{d\sigma}^\nu, \quad (155)$$

$$\mathcal{I}_a \cdot \mathcal{S}_b \cdot \mathcal{I}_c \equiv \mathcal{I}_a^\mu \mathcal{S}_{b\mu\nu}^c \mathcal{I}_c^\nu, \quad \mathcal{I}_a \cdot \mathcal{S}_b \cdot \mathcal{S}_c \cdot \mathcal{I}_d \equiv \mathcal{I}_a^\mu \mathcal{S}_{b\nu}^\mu \mathcal{S}_{c\lambda}^\nu \mathcal{I}_d^\lambda, \quad (156)$$

$$(\mathcal{S}_a \cdot \mathcal{I}_b)^\mu = \mathcal{S}_{a\nu}^\mu \mathcal{I}_b^\nu, \quad (\mathcal{S}_a \cdot \mathcal{S}_b \cdot \mathcal{I}_c)^\mu = \mathcal{S}_{a\nu}^\mu \mathcal{S}_{b\sigma}^\nu \mathcal{I}_c^\sigma, \quad (\mathcal{S}_a \cdot \mathcal{S}_b \cdot \mathcal{S}_c \cdot \mathcal{I}_d)^\mu = \mathcal{S}_{a\nu}^\mu \mathcal{S}_{b\sigma}^\nu \mathcal{S}_{c\lambda}^\sigma \mathcal{I}_d^\lambda. \quad (157)$$

2. Intensity of the collision term

The intensity part of the collision term takes the general form

$$I_{C[A]} = \frac{2^9}{2E_a} g^2 \mathcal{K} \left[\mathcal{T}_I(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \mathbf{C}, \mathbf{D}) - \mathcal{T}_I(\mathbf{A}, \mathbf{B}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right], \quad (158)$$

where the first contribution corresponds to scattering out processes and the second one to scattering in processes. The Kernel of the collision is common to both types of processes and reads

$$\begin{aligned} \mathcal{T}_I(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) \equiv & \left\{ \sum_{r=\pm} (\epsilon_r^{ac})^2 (\epsilon_r^{bd})^2 (\mathcal{I}_a^r \cdot \mathcal{I}_b^r) (\mathcal{I}_c^r \cdot \mathcal{I}_d^r) + \sum_{r=\pm} (\epsilon_r^{ac})^2 (\epsilon_{-r}^{bd})^2 (\mathcal{I}_a^r \cdot \mathcal{I}_d^{-r}) (\mathcal{I}_c^r \cdot \mathcal{I}_b^{-r}) \right. \\ & + \sum_{r=\pm} (\epsilon_r^{ac})^2 \epsilon_+^{bd} \epsilon_-^{bd} \left(+ \frac{1}{2} (\mathcal{I}_a^r \cdot \mathcal{I}_c^r) [\mathcal{S}_b \cdot \mathcal{S}_d] - (\mathcal{I}_a^r \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_c^r) - (\mathcal{I}_c^r \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_a^r) \right. \\ & \quad \left. + (\mathcal{I}_a^r \cdot \mathcal{S}_d \cdot \mathcal{I}_c^r) \mathcal{J}_b + (\mathcal{I}_c^r \cdot \mathcal{S}_b \cdot \mathcal{I}_a^r) \mathcal{J}_d + \mathcal{J}_b \mathcal{J}_d (\mathcal{I}_a^r \cdot \mathcal{I}_c^r) \right) \\ & + \sum_{r=\pm} (\epsilon_r^{bd})^2 \epsilon_+^{ac} \epsilon_-^{ac} \left(+ \frac{1}{2} (\mathcal{I}_b^r \cdot \mathcal{I}_d^r) [\mathcal{S}_a \cdot \mathcal{S}_c] - (\mathcal{I}_b^r \cdot \mathcal{S}_a \cdot \mathcal{S}_c \cdot \mathcal{I}_d^r) - (\mathcal{I}_d^r \cdot \mathcal{S}_a \cdot \mathcal{S}_c \cdot \mathcal{I}_b^r) \right. \\ & \quad \left. + (\mathcal{I}_b^r \cdot \mathcal{S}_c \cdot \mathcal{I}_d^r) \mathcal{J}_a + (\mathcal{I}_d^r \cdot \mathcal{S}_a \cdot \mathcal{I}_b^r) \mathcal{J}_c + \mathcal{J}_a \mathcal{J}_c (\mathcal{I}_b^r \cdot \mathcal{I}_d^r) \right) \\ & + \epsilon_-^{ac} \epsilon_+^{ac} \epsilon_-^{bd} \epsilon_+^{bd} \left[-[\mathcal{S}_a \cdot \mathcal{S}_c] [\mathcal{S}_b \cdot \mathcal{S}_d] + 2[\mathcal{S}_a \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{S}_c] + 2[\mathcal{S}_a \cdot \mathcal{S}_d \cdot \mathcal{S}_b \cdot \mathcal{S}_c] \right. \\ & \quad \left. - [\mathcal{S}_a \cdot \mathcal{S}_b] \mathcal{J}_c \mathcal{J}_d - [\mathcal{S}_c \cdot \mathcal{S}_d] \mathcal{J}_a \mathcal{J}_b + [\mathcal{S}_a \cdot \mathcal{S}_d] \mathcal{J}_b \mathcal{J}_c + [\mathcal{S}_b \cdot \mathcal{S}_c] \mathcal{J}_a \mathcal{J}_d + 4\mathcal{J}_a \mathcal{J}_b \mathcal{J}_c \mathcal{J}_d \right] \left. \right\}. \end{aligned} \quad (159)$$

It is rather involved since we have considered general currents (127) which include both left and right chiral coupling and we have allowed all species to be polarized. However we see immediately that if there are only left chiral couplings, only the first term in this expression survives.

3. Polarisation of the collision term

The polarisation part of the collision term must be transverse. Let us introduce the projector operator

$$H^\mu{}_\nu \equiv \delta^\mu{}_\nu + m^{-2} p^\mu p_\nu = S^\mu S_\nu + \epsilon_+^\mu \epsilon_{-\nu} + \epsilon_-^\mu \epsilon_{+\nu}. \quad (160)$$

The polarisation part of the collision term takes the general form

$$\mathcal{Q}_{C[A]}^\mu = \frac{2^9}{2E_a} g^2 \mathcal{K} (H_a)^\mu{}_\nu \left[\mathcal{T}_Q^\nu(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \mathbf{C}, \mathbf{D}) - \mathcal{T}_Q^\nu(\mathbf{A}, \mathbf{B}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right], \quad (161)$$

where $(H_a)^\mu{}_\nu$ is the projector for species a , that is orthogonally to p_a , and where the Kernel for polarisation is

$$\begin{aligned} \mathcal{T}_Q^\nu(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) \equiv & \left\{ \sum_{r=\pm} (\epsilon_r^{ac})^2 (\epsilon_r^{bd})^2 \left[+ (\mathcal{S}_a^r \cdot \mathcal{I}_b^r)^\nu (\mathcal{I}_c^r \cdot \mathcal{I}_d^r) \right] + \sum_{r=\pm} (\epsilon_r^{ac})^2 (\epsilon_{-r}^{bd})^2 \left[+ (\mathcal{S}_a^r \cdot \mathcal{I}_d^{-r})^\nu (\mathcal{I}_c^r \cdot \mathcal{I}_b^{-r}) \right] \right. \\ & + \sum_{r=\pm} \epsilon_-^{ac} \epsilon_+^{ac} (\epsilon_r^{bd})^2 \left[(\mathcal{I}_a^0 \cdot \mathcal{S}_c \cdot \mathcal{I}_b^r) \mathcal{I}_d^\nu + (\mathcal{I}_a^0 \cdot \mathcal{S}_c \cdot \mathcal{I}_d^r) \mathcal{I}_b^\nu + (\mathcal{I}_b^r \cdot \mathcal{I}_d^r) (\mathcal{S}_c \cdot \mathcal{I}_a^0)^\nu - (\mathcal{I}_b^r \cdot \mathcal{I}_a^0) (\mathcal{S}_c \cdot \mathcal{I}_d^r)^\nu \right. \\ & \quad \left. - (\mathcal{I}_d^r \cdot \mathcal{I}_a^0) (\mathcal{S}_c \cdot \mathcal{I}_b^r)^\nu - (\mathcal{I}_b^r \cdot \mathcal{I}_a^0) \mathcal{J}_c \mathcal{I}_d^\nu + (\mathcal{I}_d^r \cdot \mathcal{I}_a^0) \mathcal{J}_c \mathcal{I}_b^\nu + (\mathcal{I}_b^r \cdot \mathcal{S}_c \cdot \mathcal{I}_d^r) \mathcal{I}_a^{-r\nu} \right] \\ & + \sum_{r=\pm} \epsilon_-^{bd} \epsilon_+^{bd} (\epsilon_r^{ac})^2 \left[+ (\mathcal{S}_a^r \cdot \mathcal{S}_b^{-r} \cdot \mathcal{S}_d^r \cdot \mathcal{I}_c^r)^\nu + (\mathcal{S}_a^r \cdot \mathcal{S}_d \cdot \mathcal{S}_b \cdot \mathcal{I}_c^r)^\nu - \frac{1}{2} [\mathcal{S}_b \cdot \mathcal{S}_d] (\mathcal{S}_a^r \cdot \mathcal{I}_c^r)^\nu \right] \\ & + \epsilon_-^{ac} \epsilon_+^{ac} \epsilon_-^{bd} \epsilon_+^{bd} \left[+ 2(\mathcal{S}_d \cdot \mathcal{S}_b \cdot \mathcal{S}_c \cdot \mathcal{I}_a^0)^\nu + 2(\mathcal{S}_c \cdot \mathcal{S}_d \cdot \mathcal{S}_b \cdot \mathcal{I}_a^0)^\nu + 2(\mathcal{S}_c \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_a^0)^\nu \right. \\ & \quad + 2(\mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{S}_c \cdot \mathcal{I}_a^0)^\nu - 2[\mathcal{S}_b \cdot \mathcal{S}_d] (\mathcal{S}_c \cdot \mathcal{I}_a^0)^\nu + M_a \mathcal{Q}_a^\nu \mathcal{J}_b [\mathcal{S}_c \cdot \mathcal{S}_d] - M_a \mathcal{Q}_a^\nu \mathcal{J}_d [\mathcal{S}_b \cdot \mathcal{S}_c] \\ & \quad \left. + 4M_a \mathcal{Q}_a^\nu \mathcal{J}_b \mathcal{J}_c \mathcal{J}_d - 2\mathcal{J}_c \mathcal{J}_d (\mathcal{S}_b \cdot \mathcal{I}_a^0)^\nu + 2\mathcal{J}_c \mathcal{J}_b (\mathcal{S}_d \cdot \mathcal{I}_a^0)^\nu \right] \left. \right\}. \end{aligned} \quad (162)$$

4. Particular case of a massless species a

Expression (162) is valid also for massless case but in that case it describes only the linear polarisation part. Hence, the projector $(H_a)^\mu_\nu$ must be replaced by the screen projector $(\mathcal{H}_a)^\mu_\nu$ [see Eq. (58)] since linear polarisation is orthogonal to both p^μ and u^μ . That is the polarisation part of the collision term in the case the species a is massless takes the form

$$\mathcal{T}_Q^\nu(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) = \mathcal{T}_Q^\nu(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) + \mathcal{T}_V(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})S^\mu, \quad (163)$$

and $\mathcal{T}_Q^\nu(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ is formally equal to the rhs of Eq. (162) up to this change of projector. In the massless case, the circular polarisation part of the collision term has the structure

$$V_{C[A]} = \frac{2^9}{2E_a} g^2 \mathcal{K} \left[\mathcal{T}_V(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \mathbf{C}, \mathbf{D}) - \mathcal{T}_V(\mathbf{A}, \mathbf{B}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right], \quad (164)$$

and the Kernel for circular polarisation is

$$\begin{aligned} \lambda_a \mathcal{T}_V(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) \equiv & \left\{ \sum_{r=\pm} r (\epsilon_r^{ac})^2 (\epsilon_r^{bd})^2 (\mathcal{I}_a^r \cdot \mathcal{I}_b^r) (\mathcal{I}_c^r \cdot \mathcal{I}_d^r) + \sum_{r=\pm} r (\epsilon_r^{ac})^2 (\epsilon_{-r}^{bd})^2 (\mathcal{I}_a^r \cdot \mathcal{I}_d^{-r}) (\mathcal{I}_c^r \cdot \mathcal{I}_b^{-r}) \right. \\ & + \sum_{r=\pm} r (\epsilon_r^{ac})^2 \epsilon_+^{bd} \epsilon_-^{bd} \left[+ \frac{1}{2} (\mathcal{I}_a^r \cdot \mathcal{I}_c^r) [\mathcal{S}_b \cdot \mathcal{S}_d] - (\mathcal{I}_a^r \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_c^r) - (\mathcal{I}_c^r \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_a^r) \right. \\ & \left. \left. + (\mathcal{I}_a^r \cdot \mathcal{S}_d \cdot \mathcal{I}_c^r) \mathcal{J}_b + (\mathcal{I}_c^r \cdot \mathcal{S}_b \cdot \mathcal{I}_a^r) \mathcal{J}_d + \mathcal{J}_b \mathcal{J}_d (\mathcal{I}_a^r \cdot \mathcal{I}_c^r) \right] \right\}. \end{aligned}$$

It is possible to check that the purely circular polarisation part in the massive case, obtained from the projection along the helicity vector (40) as $S_\mu \mathcal{Q}_{C[A]}^\mu$ tends to $V_{C[A]}$ in the massless limit, showing the consistency of the expressions obtained.

E. Crossing symmetries and charge conjugation

The effect of parity and charge conjugation has been investigated in § III E. When applied on the currents which enter the interaction Hamiltonian (126), the transformation properties are found to be

$$\mathcal{C} J_{ac}^\mu(\epsilon_-^{ac}, \epsilon_+^{ac}) \mathcal{C} = J_{ac}^{\mu*}(\epsilon_+^{ac}, \epsilon_-^{ac}) \quad (165a)$$

$$\mathcal{P} J_{ac}^\mu(\epsilon_-^{ac}, \epsilon_+^{ac}) \mathcal{P} = (-1)^\mu J_{ac}^\mu(\epsilon_+^{ac}, \epsilon_-^{ac}), \quad (165b)$$

where we used the compact notation (77). To be explicit both transformations interchange the role of ϵ_-^{ac} and ϵ_+^{ac} . The additional effect of parity is not relevant because in the general Hamiltonian currents are coupled together and so $((-1)^\mu)^2 = 1$. The additional effect of charge conjugation is to add a complex conjugation. When there are no CP violating coupling matrices, such as CKM or PMNS matrices with complex phases, then this has also no effect. Combining both P and C transformations leaves thus the Hamiltonian (126) invariant. CP invariance means that the collision term for the reaction $\bar{a} + \bar{b} \leftrightarrow \bar{c} + \bar{d}$ can be either computed by applying the charge conjugation operator, or by applying the parity operation which amounts to a simple interchange of ϵ_-^{ac} and ϵ_+^{ac} .

When applying the charge conjugation operation on the collision term (146), it amounts simply to considering that all quantities built from the distribution functions (such as \mathcal{I}_μ^\pm , $S_{\mu\nu}$ etc...) now should refer to the antiparticle species. This means that the collision Kernels for the reaction $\bar{a} + \bar{b} \leftrightarrow \bar{c} + \bar{d}$ are exactly the same except that the covariant components of the antiparticles now enter its definition. That is for the collision term of species \bar{a} due to the process $\bar{a} + \bar{b} \leftrightarrow \bar{c} + \bar{d}$, the Kernel for the intensity part of the collision is given by

$$\mathcal{T}_I(\overline{\mathbf{A}}, \overline{\mathbf{B}}, \overline{\mathbf{C}}, \overline{\mathbf{D}}). \quad (166)$$

And given the CP symmetry, it can be checked explicitly that this is exactly equivalent to the interchange $\epsilon_-^{ac} \leftrightarrow \epsilon_+^{ac}$ and $\epsilon_-^{bd} \leftrightarrow \epsilon_+^{bd}$ in $\mathcal{T}_I(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$. For instance a contribution of the form

$$\sum_r (\epsilon_r^{ac})^2 (\epsilon_r^{bd})^2 (\mathcal{I}_a^r \cdot \mathcal{I}_b^r) (\mathcal{I}_c^r \cdot \mathcal{I}_d^r) \quad (167)$$

when considered for antiparticles is equivalent to the operation $\epsilon_-^{ac} \leftrightarrow \epsilon_+^{ac}$ and $\epsilon_-^{bd} \leftrightarrow \epsilon_+^{bd}$ because \mathcal{I}_μ^\pm for a particle is equal to \mathcal{I}_μ^\mp of the corresponding antiparticle.

Now that we have computed the collision term for $a + b \leftrightarrow c + d$ and deduced in the most straightforward manner the one associated to $\bar{a} + \bar{b} \leftrightarrow \bar{c} + \bar{d}$, we can obtain all other related reactions by crossing symmetry. Let us for instance consider the processes $a \leftrightarrow \bar{b} + c + d$. The part of the interaction Hamiltonian responsible for this process, $H_I^{a \leftrightarrow \bar{b} + c + d}$ can be deduced from $H_I^{a + b \leftrightarrow c + d}$ given in Eq. (138) by the replacement

$$b_s \rightarrow \bar{b}_s^\dagger, \quad b_s^\dagger \rightarrow \bar{b}_s, \quad u_s \rightarrow v_s, \quad \bar{u}_s \rightarrow \bar{v}_s. \quad (168)$$

From Eqs. (B2), we see that for the species b , the Pauli blocking operator of the particle species b , \widehat{B}_{rs} , (resp the number operator B_{rs}) is replaced by number operator of the antiparticle species \bar{b} , \overline{B}_{rs} , (resp. the Pauli blocking operator $\widehat{\overline{B}}_{rs}$), which was expected since we have changed an initial state for a final state. The change from a particle to an antiparticle is also expected and consistent with the contraction of the operators which is now made with v_s and \bar{v}_s instead of u_s and \bar{u}_s when defining the associated spinor valued operators as in Eqs. (38). For instance the intensity part due to the process $a + \bar{c} \leftrightarrow \bar{b} + d$ on the species a is

$$I_{C[A]} = \frac{2^9}{2E_a} g^2 \mathcal{K} \left[\mathcal{T}_I(\widehat{A}, \overline{B}, \widehat{C}, D) - \mathcal{T}_I(A, \widehat{B}, \overline{C}, \widehat{D}) \right], \quad (169)$$

where we made clear with the barred notation that the covariant quantities related to the species \bar{b} and \bar{c} are those of antiparticles. These rules for crossing symmetry apply as well for the polarisation part of the collision term.

F. Structure of the collision Kernels

Before applying these general results to particular examples corresponding to physical situations, let us comment on the general structure of the collision terms.

- The intensity Kernel satisfies the property

$$\mathcal{T}_I(A, B, C, D) = \mathcal{T}_I(C, D, A, B). \quad (170)$$

Since the simple interaction (126) has a CP symmetry, then it must also have a T (time-reversal) symmetry to ensure CPT symmetry and this is reflected by this relation. This property could also have been found from Eq. (147).

- In the case where the species a whose collision term is considered is massless, the Kernel for the circular polarisation (165) does not involve couplings of the left and right chiral coupling constants ϵ_-^{ac} and ϵ_+^{ac} , since there are no terms proportional to $\epsilon_-^{ac} \epsilon_+^{ac}$. Furthermore, if no species is initially polarized (circularly or linearly), then if the chiral coupling constants are equal ($\epsilon_-^{ac} = \epsilon_+^{ac}$ and $\epsilon_-^{bd} = \epsilon_+^{bd}$), the Kernel for circular polarisation vanishes. A departure from a purely unpolarized state must be due to a difference between the left and right chiral couplings in one of the currents.
- In the massless case it is also instructive to consider the particular case when one of the chiral couplings constants ϵ_-^{ac} or ϵ_+^{ac} vanishes, corresponding to either a purely left or purely right chiral coupling in the current of the ac pair. In that case, we can focus on the purely left or purely right helicity part of the collision term by considering

$$\begin{aligned} \mathcal{T}_I(A, B, C, D) \pm \lambda_a \mathcal{T}_V(A, B, C, D) &= (\epsilon_\pm^{ac})^2 (\epsilon_\pm^{bd})^2 (\mathcal{I}_a^\pm \cdot \mathcal{I}_b^\pm) (\mathcal{I}_c^\pm \cdot \mathcal{I}_d^\pm) + (\epsilon_\pm^{ac})^2 (\epsilon_\mp^{bd})^2 (\mathcal{I}_a^\pm \cdot \mathcal{I}_d^\mp) (\mathcal{I}_c^\pm \cdot \mathcal{I}_b^\mp) \\ &+ (\epsilon_\pm^{ac})^2 \epsilon_\mp^{bd} \epsilon_\mp^{bd} \left[+ \frac{1}{2} (\mathcal{I}_a^\pm \cdot \mathcal{I}_c^\pm) (\mathcal{S}_b \cdot \mathcal{S}_d) - (\mathcal{I}_a^\pm \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_c^\pm) - (\mathcal{I}_c^\pm \cdot \mathcal{S}_b \cdot \mathcal{S}_d \cdot \mathcal{I}_a^\pm) \right. \\ &\left. + (\mathcal{I}_a^\pm \cdot \mathcal{S}_d \cdot \mathcal{I}_c^\pm) \mathcal{J}_b + (\mathcal{I}_c^\pm \cdot \mathcal{S}_b \cdot \mathcal{I}_a^\pm) \mathcal{J}_d + \mathcal{J}_b \mathcal{J}_d (\mathcal{I}_a^\pm \cdot \mathcal{I}_c^\pm) \right]. \end{aligned} \quad (171)$$

If the species a is a particle $\lambda_a = 1$ (resp. an antiparticle $\lambda_a = -1$), and if $\epsilon_+^{ac} = 0$ (resp. $\epsilon_-^{ac} = 0$), then this expression means that only the left helicities $I_a - V_a$ are affected by the collision term (resp. only the right helicities $I_a + V_a$). If the species a is also not polarized linearly, then the linear polarisation part of the collision term vanishes as well under these assumptions. Hence we recover that in the massless case, for interactions which have a definite chirality for a given species, then if the distribution of particles is maximally circularly polarized ($I = V$ or $I = -V$), that is such that only one type of helicity is present and corresponds to the chirality of the interaction, then it remains so even after interactions. The other helicity states can only be sourced if the interactions are not purely chiral (both $\epsilon_\pm^{ac} \neq 0$) or if the particles are massive.

For instance at very high energies when we can consider that both neutrinos and electrons are massless, neutrinos are only left chiral and antineutrinos are only right chiral and they remain so because the currents with which they couple (122) and (118) are always purely left chiral, whereas for electrons both helicities are populated since the neutral currents (119) involve left and right chiral coupling constants.

- This feature is a particular case of a more general feature of the collision term. Indeed, in § VI C, we have commented on the fact that the collision term only affects one of the two spin indices in the distribution function \mathcal{A}_{rs} . This means that a state with only positive helicities \mathcal{A}_{++} (resp. \mathcal{A}_{--}) either evolves under one collision to \mathcal{A}_{++} (resp. \mathcal{A}_{--}), \mathcal{A}_{-+} or \mathcal{A}_{+-} . And the two latter states can evolve toward \mathcal{A}_{--} (resp. \mathcal{A}_{++}) after a second collision. A distribution function which is fully circularly polarized (such that $I = \pm V$), that is which is made only of helicity states of one kind, must first interact to generate linear polarisation and then interact again to generate helicity states of the other kind. In the massless case and for the case $\epsilon_+^{ac} = 0$, given that chirality is the same thing as helicity, then there is no transition possible from \mathcal{A}_{--} to \mathcal{A}_{-+} or \mathcal{A}_{+-} and the distribution with purely left helicities remains so even after collisions.

VII. APPLICATION TO STANDARD REACTIONS

In practice the general Kernels for the collision term obtained in the previous section are far too general. Indeed, we shall consider cases in which only one species is polarized or could be polarized by the collisions. For instance when considering electrons, apart in certain circumstances, it is not useful to describe their polarisation state since electromagnetic interactions will contribute to erase any polarisation. Furthermore at high energies compared to the masses of neutrinos, it is reasonable to approximate the neutrinos as being massless. Under these two types of simplifications (absence of polarisation or of mass for some species) the collision Kernels take simpler forms. Finally for nearly all applications of interest, some chiral coupling constants vanish.

A. Processes involving 4 bodies

1. Muon decay

The muon decay is due to the interaction between the muon (μ^-)/muon neutrino (ν_μ) charged current and the electron (e^-)/neutrino (ν) charged current. Furthermore it involves only left-chiral couplings. It thus corresponds to the case

$$a = \mu^-, \quad c = \nu_\mu, \quad b = \nu, \quad d = e^-, \quad \epsilon_+^{ac} = \epsilon_+^{bd} = 0, \quad \epsilon_-^{ac} = \epsilon_-^{bd} = 1, \quad g = \frac{G_F}{\sqrt{2}}. \quad (172)$$

We remind that for the decay reaction $a \leftrightarrow \bar{b} + c + d$, the collision term is deduced from the reaction $a + b \leftrightarrow c + d$ by crossing symmetry. For instance from Eqs. (159) and (162), we deduce that the covariant parts of the collision term are of the form

$$I_{C[A]} = \frac{2^8}{2E_a} G_F^2 \mathcal{K} \left[\mathcal{T}_I(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \mathbf{C}, \mathbf{D}) - \mathcal{T}_I(\mathbf{A}, \widehat{\mathbf{B}}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right] \quad (173a)$$

$$\mathcal{Q}_{C[A]}^\mu = \frac{2^8}{2E_a} G_F^2 \mathcal{K} (H_a)^\mu{}_\nu \left[\mathcal{T}_Q^\nu(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \mathbf{C}, \mathbf{D}) - \mathcal{T}_Q^\nu(\mathbf{A}, \widehat{\mathbf{B}}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right] \quad (173b)$$

and where it is stressed by a barred notation that the covariant quantities related to the species a , c and d are those of particles, and those for the species \bar{b} are those of antiparticles. The Kernels are the most simple ones and read

$$\mathcal{T}_I(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) = (\mathcal{I}_a^- \cdot \mathcal{I}_b^-)(\mathcal{I}_c^- \cdot \mathcal{I}_d^-) \quad (174a)$$

$$\mathcal{T}_Q^\nu(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) = (\mathcal{S}_a^- \cdot \mathcal{I}_b^-)^\nu (\mathcal{I}_c^- \cdot \mathcal{I}_d^-). \quad (174b)$$

If we neglect the three body reactions, we need only to consider the scattering out processes which correspond to the muon decay. Furthermore if all species are unpolarized, and if the density of species b, c, d is low enough such that we can neglect their Pauli blocking effects, then the intensity part of the collision term becomes

$$I_{C[A]}(p) = -\frac{32}{E_a} G_F^2 I_a(p) \int [dp_b][dp_c][dp_d] (2\pi)^4 \delta^{(4)}(p_d + p_c - p_b - p) (p \cdot p_b) (p_c \cdot p_d). \quad (175)$$

As a check, the muon lifetime is recovered from this collision term evaluated at null spatial momentum of ($\mathbf{p} = 0$) thanks to the definition $I_{C[A]}(\mathbf{p} = 0) = dI_a/dt(\mathbf{p} = 0) \equiv -\Gamma_a I_a(\mathbf{p} = 0)$. Using $p \cdot p_b = -m_a E_b$, the muon lifetime is obtained as

$$\Gamma_a = 32G_F^2 \int [dp_b][dp_c][dp_d] (2\pi)^4 \delta^{(3)}(\mathbf{p}_d + \mathbf{p}_c - \mathbf{p}_b) \delta^{(1)}(E_d + E_c - E_b - m_a) E_b (-p_c \cdot p_d), \quad (176)$$

and this is exactly the expression that would be obtained from the Fermi golden rule.

Finally let us comment on the structure of the polarisation Kernel (174b) corresponding to the processes $a+b \leftrightarrow c+d$. Indeed, since the currents coupled are purely left chiral, the structure of the collision term is much more transparent than the general case.

- The Kernel (174b) has to be projected in Eq. (173b) to form the polarisation part of the collision. From the definition (154) for $\mathcal{S}_{\mu\nu}^-$ we see that the term involving $\mathcal{S}_{\mu\nu}$ cannot contribute if muons are initially not polarized. Even if they are linearly polarized, this contribution does not enhance polarisation since either the projection acts on p_a and thus it cancels, or it acts on \mathcal{Q}_a^μ , but if it is a scattering out process this reduces polarisation, and if it is a scattering in process, then this is a Pauli blocking form and also reduces polarisation. Hence one can replace $(\mathcal{S}_a^- \cdot \mathcal{I}_b^-)^\nu$ by $+M_a \mathcal{I}_a \mathcal{I}_b^{-\nu}$. The first conclusion we draw is that polarisation can be generated for muons only because they have a mass. If they were massless, no polarisation could be generated. Furthermore, in the case where species b, c, d are unpolarized, then $\mathcal{I}_b^{-\nu} = \mathcal{I}_b p_b^\nu$ and so the polarisation part of the collision term is proportional to $(H_a)^\mu{}_\nu p_b^\nu$, that is to the transverse part (with respect to p_a^μ) of the momentum of species b .
- Focusing on the linear polarisation part of the collision term, it is sufficient to consider the spatial components. Neglecting the Pauli blocking effects, the part due to scattering in processes for the purely linear polarisation part of the collision term in the reaction $a + b \leftrightarrow c + d$ has the structure

$$Q_{C[A]}^i(p) \propto m_a \int \frac{[dp_c][dp_d]}{E_b} \delta(E_c + E_d - E - E_b) (\delta_j^i - \hat{p}^i \hat{p}_j) (p_c^j + p_d^j) I_c(\mathbf{p}_c) I_d(\mathbf{p}_d) (p_c \cdot p_d). \quad (177)$$

The integrals on momenta $[dp_c][dp_d]$ can be separated into direction and magnitude. One integral on magnitude (either $|p_c|$ or $|p_d|$) can be used to remove the Dirac delta function on energies. Let us for simplicity ignore the directional dependence in the factors $(p_c \cdot p_d)$ and $1/E_b$. The integral on the direction \hat{p}_c of the factor $p_c^j I_c(\mathbf{p}_c)$ selects by definition the dipolar contribution ($\ell = 1$ when expanded in spherical harmonics) of species c , and similarly the integral on \hat{p}_d of the factor $p_d^j I_d(\mathbf{p}_d)$ selects the dipolar part of species d . Furthermore the projection operator $(\delta_j^i - \hat{p}^i \hat{p}_j)$ ensures that the linear polarisation generated has also exactly a dipolar structure (we remind from § IV B that the angular expansion is performed with spin- ± 1 spherical harmonics). The reasoning is more complex when correctly taking into account the directional dependence in $(p_c \cdot p_d)$ and $1/E_b$ but the conclusion remains the same, and it is that we need a dipolar structure in the initial species c or d to create polarisation in the massive particle a . This could also have been guessed from the angular decompositions of § IV B since linear polarisation is decomposed in spin- ± 1 spherical harmonics, and these cannot have a monopole ($\ell = m = 0$).

- However, nothing prevents circular polarisation to develop, and it does indeed so because the couplings are purely left chiral, so only pure helicity states of one kind are affected by collisions. The circular part of the collision term is extracted by projection of (173b) on the helicity vector S^μ . In particular, if we could neglect the masses of muons, for instance at very high energies, then we would find that only $I - V$ is sourced, that is only purely left helicity states are affected by collisions, as discussed already in the general case (171). Note that since circular polarisation is expanded in standard spherical harmonics, nothing prevents purely monopolar distributions (for instance as those of homogeneous cosmology) to generate circular polarisation, even though it does not generate linear polarisation. A different observer, that is an observer boosted with respect to the one which sees only purely monopolar distributions, would see distributions with a dipolar structure, and thus would see that the collision term produces linear polarisation. There is no contradiction because what appears as purely circular polarisation for an observer, appears as both circular and linear polarisation for the boosted one. Only the total polarisation Q^μ is observer independent.
- In the massless case, both circular and linear polarisation are observer independent, but as we have seen, linear polarisation is not generated, and for all observers only intensity and circular polarisation are affected by collisions.

2. Neutron decay

The neutron decay is due to the coupling of the neutron (n)/proton (p) charged current and the neutrino/electron charged current. While the latter is purely left chiral, the former has both chiral couplings due to the effective constant

g_A defined in Eq. (123). Hence we must consider the case

$$a = n, \quad c = p, \quad b = \nu, \quad d = e^-, \quad \epsilon_+^{ac} = \frac{1-g_A}{2}, \quad \epsilon_+^{bd} = 0, \quad \epsilon_-^{ac} = \frac{1+g_A}{2}, \quad \epsilon_-^{bd} = 1, \quad g = \frac{G_F \cos \theta_C}{\sqrt{2}}. \quad (178)$$

The neutron decay is deduced by crossing symmetry from the reaction $n + \nu \leftrightarrow p + e^-$ (and so is the reaction $n + e^+ \leftrightarrow p + \bar{\nu}$). As in the muon decay, the covariant parts of the collision term are of the form (173), but now the Kernels are slightly more complicated due to the non-vanishing ϵ_+^{ac} . For instance the intensity part of the collision term for the neutron decay takes the same form as (173a), with an extra CKM angle factor $(\cos \theta_C)^2$, but with the Kernel

$$\begin{aligned} \mathcal{T}_I(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) &= \left(\frac{1+g_A}{2}\right)^2 (\mathcal{I}_a^- \cdot \mathcal{I}_b^-)(\mathcal{I}_c^- \cdot \mathcal{I}_d^-) + \left(\frac{1-g_A}{2}\right)^2 (\mathcal{I}_a^+ \cdot \mathcal{I}_d^-)(\mathcal{I}_c^+ \cdot \mathcal{I}_b^-) \\ &+ \left(\frac{1-g_A^2}{4}\right) \left(+\frac{1}{2}(\mathcal{I}_b^- \cdot \mathcal{I}_d^-)[\mathcal{S}_a \cdot \mathcal{S}_c] - (\mathcal{I}_b^- \cdot \mathcal{S}_a \cdot \mathcal{S}_c \cdot \mathcal{I}_d^-) - (\mathcal{I}_d^- \cdot \mathcal{S}_a \cdot \mathcal{S}_c \cdot \mathcal{I}_b^-) \right. \\ &\quad \left. + (\mathcal{I}_b^- \cdot \mathcal{S}_c \cdot \mathcal{I}_d^-)\mathcal{J}_a + (\mathcal{I}_d^- \cdot \mathcal{S}_a \cdot \mathcal{I}_b^-)\mathcal{J}_c + \mathcal{J}_a\mathcal{J}_c(\mathcal{I}_b^- \cdot \mathcal{I}_d^-) \right). \end{aligned} \quad (179)$$

If no species is polarized, the Kernel takes the simpler form

$$\begin{aligned} \mathcal{T}_I(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) &= \mathcal{I}_a \mathcal{I}_b \mathcal{I}_c \mathcal{I}_d \left[\left(\frac{1+g_A}{2}\right)^2 (p_a \cdot p_b)(p_c \cdot p_d) + \left(\frac{1-g_A}{2}\right)^2 (p_a \cdot p_d)(p_b \cdot p_c) \right. \\ &\quad \left. - \left(\frac{g_A^2-1}{4}\right) M_a M_c (p_b \cdot p_d) \right]. \end{aligned} \quad (180)$$

This form of the Kernel matches Eq. (3.25) of Ref. [23]. The life-time of the neutron at zero temperature (that is when we can neglect the Pauli blocking factors), is obtained from the same method as for the lifetime of the muon (176) and we get

$$\begin{aligned} \Gamma_a &= -32(G_F \cos \theta_C)^2 \int [dp_b][dp_c][dp_d] (2\pi)^4 \delta^{(3)}(\mathbf{p}_d + \mathbf{p}_c - \mathbf{p}_b) \delta^{(1)}(E_d + E_c - E_b - m_a) \\ &\times \left[\left(\frac{1+g_A}{2}\right)^2 E_b (p_c \cdot p_d) + \left(\frac{1-g_A}{2}\right)^2 E_d (p_b \cdot p_c) + \left(\frac{g_A^2-1}{4}\right) m_c (p_b \cdot p_d) \right]. \end{aligned} \quad (181)$$

It is customary to evaluate this decay rate in the infinite mass limit for the protons and neutrons, that is using the rule

$$E_c = m_c, \quad p_c \cdot p_d = -m_c E_d, \quad p_b \cdot p_c = -m_c E_b, \quad p_b \cdot p_d = -E_b E_d + \mathbf{p}_b \cdot \mathbf{p}_d. \quad (182)$$

Integrating over the spatial momentum of the protons we obtain

$$\Gamma_a \simeq \frac{(G_F \cos \theta_C)^2}{(2\pi)^5} \int d^3 \mathbf{p}_b d^3 \mathbf{p}_d \delta^{(1)}(E_d + m_c - E_b - m_a) \left[(1 + 3g_A^2) + (1 - g_A^2) \frac{\mathbf{p}_b \cdot \mathbf{p}_d}{E_b E_d} \right] \quad (183a)$$

$$\simeq (1 + 3g_A^2) \frac{(G_F \cos \theta_C)^2}{2\pi^3} \int |\mathbf{p}_d| E_d (E_d + m_c - m_a)^2 dE_d, \quad (183b)$$

where in the last step we have used that an integral on spatial momenta directions of $\mathbf{p}_p \cdot \mathbf{p}_d$ vanishes since it is odd, and we have considered the neutrinos as being totally massless. Hence we recover the standard result for the neutron decay in the $V - A$ theory (see e.g. Refs. [11, 42–45]).

3. Big-bang nucleosynthesis reactions

We have already mentioned in § VI E that from the collision term for neutrons due to the reaction $n + \nu \leftrightarrow p + e^-$ we can deduce the neutron decay and $n + e^+ \leftrightarrow p + \bar{\nu}$ thanks to crossing symmetry. To be complete, let us illustrate the procedure to get all the collision terms for these four species, focusing on a few examples.

- In order to get the collision term for neutrinos, we must consider the reaction $\nu + n \leftrightarrow e^- + p$, that is we must consider

$$a = \nu, \quad c = e^-, \quad b = n, \quad d = p, \quad \epsilon_+^{bd} = \frac{1-g_A}{2}, \quad \epsilon_+^{ac} = 0, \quad \epsilon_-^{bd} = \frac{1+g_A}{2}, \quad \epsilon_-^{ac} = 1, \quad g = \frac{G_F \cos \theta_C}{\sqrt{2}}. \quad (184)$$

- In order to get the collision term for electrons we must consider the reaction $e^- + p \leftrightarrow \nu + n$, that is the case

$$a = e^-, \quad c = \nu, \quad b = p, \quad d = n, \quad \epsilon_+^{bd} = \frac{1-g_A}{2}, \quad \epsilon_+^{ac} = 0, \quad \epsilon_-^{bd} = \frac{1+g_A}{2}, \quad \epsilon_-^{ac} = 1, \quad g = \frac{G_F \cos \theta_C}{\sqrt{2}}. \quad (185)$$

- In order to get the collision term for antineutrinos which is due to the reaction $\bar{\nu} + p \leftrightarrow e^+ + n$, we must consider the reaction $\bar{\nu} + \bar{n} \leftrightarrow e^+ + \bar{p}$ and deduce the former by crossing symmetry as explained in § VIE. This means that we consider the case

$$a = \bar{\nu}, \quad c = e^+, \quad b = n, \quad d = p, \quad \epsilon_+^{bd} = \frac{1-g_A}{2}, \quad \epsilon_+^{ac} = 0, \quad \epsilon_-^{bd} = \frac{1+g_A}{2}, \quad \epsilon_-^{ac} = 1, \quad g = \frac{G_F \cos \theta_C}{\sqrt{2}}. \quad (186)$$

In all these case, there is a polarisation component in the collision term. Its detailed expression is more complex than Eq. (173b) because the couplings are not purely left chiral but the general structure remains the same. Circular polarisation is generated in all cases, and linear polarisation is generated if the distribution functions of the interacting species have some dipolar distribution. Since the electron/neutrino current is still purely left chiral, then if these were the only interactions, the electrons and neutrinos would develop circular polarisation. If this is certainly the case for massless neutrinos, which develop maximal circular polarisation, this would not be the case for electrons which have both right and chiral couplings thanks to neutral current interactions, and also more importantly due to electromagnetic interactions which are far stronger in general and would erase any type of polarisation.

All the four-body reactions related to the neutron beta decay are gathered in Table 2.

Reaction	Particles names	Chiral couplings				$2^{-7} G_F^{-2} E I_{C[A]} = \mathcal{K} \times$
$n + \nu \leftrightarrow p + e^-$	$a + b \leftrightarrow c + d$	$\epsilon_-^{ac} = \frac{(1+g_A)}{2}$	$\epsilon_+^{ac} = \frac{(1-g_A)}{2}$	$\epsilon_-^{bd} = 1$	$\epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})$
$p + e^- \leftrightarrow n + \nu$	$a + b \leftrightarrow c + d$	$\epsilon_-^{ac} = \frac{(1+g_A)}{2}$	$\epsilon_+^{ac} = \frac{(1-g_A)}{2}$	$\epsilon_-^{bd} = 1$	$\epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})$
$\nu + n \leftrightarrow e^- + p$	$a + b \leftrightarrow c + d$	$\epsilon_-^{bd} = \frac{(1+g_A)}{2}$	$\epsilon_+^{bd} = \frac{(1-g_A)}{2}$	$\epsilon_-^{ac} = 1$	$\epsilon_+^{ac} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})$
$e^- + p \leftrightarrow \nu + n$	$a + b \leftrightarrow c + d$	$\epsilon_-^{bd} = \frac{(1+g_A)}{2}$	$\epsilon_+^{bd} = \frac{(1-g_A)}{2}$	$\epsilon_-^{ac} = 1$	$\epsilon_+^{ac} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})$
$n + e^+ \leftrightarrow p + \bar{\nu}$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{ac} = \frac{(1+g_A)}{2}$	$\epsilon_+^{ac} = \frac{(1-g_A)}{2}$	$\epsilon_-^{bd} = 1$	$\epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \widehat{D})$
$p + \bar{\nu} \leftrightarrow n + e^+$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{ac} = \frac{(1+g_A)}{2}$	$\epsilon_+^{ac} = \frac{(1-g_A)}{2}$	$\epsilon_-^{bd} = 1$	$\epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \widehat{D})$
$e^+ + n \leftrightarrow \bar{\nu} + p$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{bd} = \frac{(1+g_A)}{2}$	$\epsilon_+^{bd} = \frac{(1-g_A)}{2}$	$\epsilon_-^{ac} = 1$	$\epsilon_+^{ac} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \widehat{D})$
$\bar{\nu} + p \leftrightarrow e^+ + n$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{bd} = \frac{(1+g_A)}{2}$	$\epsilon_+^{bd} = \frac{(1-g_A)}{2}$	$\epsilon_-^{ac} = 1$	$\epsilon_+^{ac} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \widehat{D})$
$n \leftrightarrow \bar{\nu} + p + e^-$	$a \leftrightarrow \bar{b} + c + d$	$\epsilon_-^{ac} = \frac{(1+g_A)}{2}$	$\epsilon_+^{ac} = \frac{(1-g_A)}{2}$	$\epsilon_-^{bd} = 1$	$\epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \widehat{D})$

FIG. 2: Main four-body reactions related to the neutron beta decay. The general intensity Kernel \mathcal{T}_I is given in § VID by Eq. (159). The polarisation part of the collision term $\mathcal{Q}_{C[A]}^\mu$ is obtained by using instead $H_a^{\mu\nu} \mathcal{T}_C^\nu$, where the Kernel for polarisation is given in the general case by Eq. (162). If species a is massless, one must distinguish between the purely linear polarisation part and the circular polarisation part of the collision term as explained in § VID4. In all cases, the collision term is the one of particle a or antiparticle \bar{a} . In the arguments of the Kernels, we have either particles or antiparticles operators, and in the latter cases we use a barred notation to stress that the associated covariant components (\mathcal{T}_μ^\pm , $\mathcal{S}_{\mu\nu}$ etc...) which enter the expression of the Kernel are those of antiparticles. Furthermore for final states, operators are indicated with a hatted notation, since we must take the Pauli blocking forms, that is the hatted forms, of the covariant components defined in § VID.

B. Processes involving 2 bodies

There are in general two types of 2-bodies interactions. Either reactions of thermalisation of the type $a + b \leftrightarrow a + b$ or annihilation reactions which enforce chemical equilibrium such as $a + \bar{a} \leftrightarrow b + \bar{b}$ [9]. As analyzed in § VI E, using crossing symmetry one can deduce the latter from the former. Additionally all cases can be reduced to the interaction of neutral currents with a simple Fierz reordering as in Eq. (124). The two-body process can thus be considered as special cases of the 4-body processes for which

$$c = a, \quad d = b, \quad (187)$$

and the interaction Hamiltonian due to neutral current takes the form

$$\mathcal{H}_I = -\mathcal{L}_I = -g J_\mu^{aa} J_{bb}^\mu, \quad g = 2 \frac{G_F}{\sqrt{2}}. \quad (188)$$

This extra factor 2 in the coupling constant g can be viewed as the contribution of the complex conjugated terms in Eq. (126). The general derivation of the collision term follows the same general steps as in § VI C. However Eq. (141) is modified since operators of species c and a do not trivially commute as they are the same species and similarly for species b and d . We must instead use

$$\langle [\mathcal{M}, [\mathcal{M}, A_{ss'}]] \rangle \equiv M_{\alpha\beta \rightarrow \gamma'\delta'}^* M_{\alpha'\beta' \rightarrow \gamma\delta} \langle [b_\beta^\dagger b_{\delta'} a_\alpha^\dagger a_{\gamma'} + b_\delta^\dagger b_{\beta'} [a_\gamma^\dagger a_{\alpha'} + a_s^\dagger a_{s'}]] \rangle, \quad (189)$$

$$\begin{aligned} \langle [b_\beta^\dagger b_{\delta'} a_\alpha^\dagger a_{\gamma'} + b_\delta^\dagger b_{\beta'} [a_\gamma^\dagger a_{\alpha'} + a_s^\dagger a_{s'}]] \rangle &= \langle [B_{\beta\delta'} A_{\alpha\gamma'}, B_{\delta\beta'} (A_{\gamma s'} \underline{\delta}_{s\alpha'} - A_{s\alpha'} \underline{\delta}_{\gamma s'})] \rangle \\ &= \langle \left\{ (B_{\beta\beta'} \widehat{B}_{\delta\delta'} + B_{\beta\delta'} B_{\delta\beta'}) \left[(A_{\alpha s'} \widehat{A}_{\gamma\gamma'} + A_{\alpha\gamma'} A_{\gamma s'}) \underline{\delta}_{s\alpha'} - (A_{\alpha\alpha'} \widehat{A}_{s\gamma'} + A_{\alpha\gamma'} A_{s\alpha'}) \underline{\delta}_{\gamma s'} \right] \right. \\ &\quad \left. - (B_{\delta\delta'} \widehat{B}_{\beta\beta'} + B_{\beta\delta'} B_{\delta\beta'}) \left[(A_{\gamma\gamma'} \widehat{A}_{\alpha s'} + A_{\alpha\gamma'} A_{\gamma s'}) \underline{\delta}_{s\alpha'} - (A_{s\gamma'} \widehat{A}_{\alpha\alpha'} + A_{\alpha\gamma'} A_{s\alpha'}) \underline{\delta}_{\gamma s'} \right] \right\} \rangle, \end{aligned} \quad (190)$$

where for the first equality we have use property (B6) and for the second line we have used the property (30). The difference lies in the fact that there are additional terms with distribution functions such as $\mathcal{B}_{\beta\delta'}$, $\widehat{\mathcal{B}}_{\delta\beta'}$ or $\mathcal{A}_{\alpha\gamma'}$, that is such that indices are not pairs $\alpha s'$, $s\alpha'$, $\beta\beta'$, $\gamma\gamma'$ or $\delta\delta'$ as in Eq. (143). However, when taking the expectation in the quantum state this introduces Dirac delta functions [from Eq. (1)], and for these types of terms it corresponds to a process where the momenta of species a and b between the initial and final state are unchanged, that is to processes where there is no momentum exchange at all. It is expected that these processes should be irrelevant and that scattering out and scattering in processes due to these terms cancel identically. We conclude that the analysis of § VI C, and the general features of the collision term we have subsequently highlighted, can be completely transposed for the two-body processes, provided we use (187). In order to give concrete example we apply it to some two-body interactions mediated by weak interactions.

1. neutrino/muon neutrino scattering

The interactions between neutrinos of different types (e.g. electronic neutrinos and muonic neutrinos) are only due to neutral currents with a pure left chiral coupling. The effect of the reaction $\nu + \nu_\mu \leftrightarrow \nu + \nu_\mu$ thus corresponds to the case

$$a = \nu, \quad b = \nu_\mu, \quad c = \nu, \quad d = \nu_\mu, \quad \epsilon_-^{ac} = \epsilon_-^{bd} = e_-^\nu = \frac{1}{2}, \quad \epsilon_+^{ac} = \epsilon_+^{bd} = 0, \quad g = 2 \frac{G_F}{\sqrt{2}}. \quad (191)$$

If we cannot neglect the masses of neutrinos, then the covariant parts of the collision term take the form

$$I_{C[A]} = \frac{2^{10}}{2E_a} G_F^2 \mathcal{K} \left[\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D}) \right] \quad (192a)$$

$$\mathcal{Q}_{C[A]}^\mu = \frac{2^{10}}{2E_a} G_F^2 \mathcal{K} (H_a)^\mu{}_\nu \left[\mathcal{T}_Q^\nu(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_Q^\nu(A, B, \widehat{C}, \widehat{D}) \right] \quad (192b)$$

where the Kernels are $1/2^4$ times those of the muon decay (174), that is

$$\mathcal{T}_I(A, B, C, D) = \frac{1}{2^4} (\mathcal{I}_a^- \cdot \mathcal{I}_b^-) (\mathcal{I}_c^- \cdot \mathcal{I}_d^-) \quad (193a)$$

$$\mathcal{T}_Q^\nu(A, B, C, D) = \frac{1}{2^4} (\mathcal{S}_a^- \cdot \mathcal{I}_b^-)^\nu (\mathcal{I}_c^- \cdot \mathcal{I}_d^-). \quad (193b)$$

If we can consider that the electronic neutrinos are massless, then we must also consider the collision term for circular polarisation since Eq. (192b) only applies to the linear polarisation part when the projector $(H_a)^\mu_\nu$ is replaced by the screen projector $(\mathcal{H}_a)^\mu_\nu$. The circular polarisation part of the collision term reads simply

$$\lambda_a V_{C[A]} = \frac{2^{10}}{2E_a} G_F^2 \mathcal{K} \left[\mathcal{T}_V(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \mathbf{C}, \mathbf{D}) - \mathcal{T}_V(\mathbf{A}, \mathbf{B}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right] \quad (194)$$

$$\mathcal{T}_V(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) = -\frac{1}{2^4} (\mathcal{I}_a^- \cdot \mathcal{I}_b^-) (\mathcal{I}_c^- \cdot \mathcal{I}_d^-). \quad (195)$$

The effect of the reaction $\nu + \bar{\nu}_\mu \leftrightarrow \nu + \bar{\nu}_\mu$, which in our general notation is $a + \bar{d} \leftrightarrow c + \bar{b}$, is obtained by a simple crossing symmetry. For instance the intensity part of the Kernel would be for that process

$$I_{C[A]} = \frac{2^{10}}{2E_a} G_F^2 \mathcal{K} \left[\mathcal{T}_I(\widehat{\mathbf{A}}, \overline{\mathbf{B}}, \mathbf{C}, \widehat{\mathbf{D}}) - \mathcal{T}_I(\mathbf{A}, \overline{\mathbf{B}}, \widehat{\mathbf{C}}, \overline{\mathbf{D}}) \right], \quad (196)$$

where we emphasized again with a barred notation that the covariant components of antiparticles are those of antiparticles.

For completeness, we must stress again that the effect of antineutrino-muonic antineutrino reactions ($\bar{\nu} + \bar{\nu}_\mu \leftrightarrow \bar{\nu} + \bar{\nu}_\mu$) on antineutrinos is obtained by charge conjugation, that is by considering the case

$$a = \bar{\nu}, \quad b = \bar{\nu}_\mu, \quad c = \bar{\nu}, \quad d = \bar{\nu}_\mu, \quad \epsilon_-^{ac} = \epsilon_-^{bd} = e_-^\nu = \frac{1}{2}, \quad \epsilon_+^{ac} = \epsilon_+^{bd} = 0, \quad g = 2 \frac{G_F}{\sqrt{2}}. \quad (197)$$

This means that the collision term takes the same form as Eqs. (192) but where all covariant components should now refer to antiparticle species. For instance the intensity part takes the form

$$I_{C[A]} = \frac{2^{10}}{2E_a} G_F^2 \mathcal{K} \left[\mathcal{T}_I(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \overline{\mathbf{C}}, \overline{\mathbf{D}}) - \mathcal{T}_I(\overline{\mathbf{A}}, \overline{\mathbf{B}}, \widehat{\mathbf{C}}, \widehat{\mathbf{D}}) \right]. \quad (198)$$

The effect of the process $\bar{\nu} + \nu_\mu \leftrightarrow \bar{\nu} + \nu_\mu$ is then obtained by a crossing symmetry as in § VI E.

2. neutrino/neutrino scattering

Neutrino-neutrino scattering ($\nu + \nu \leftrightarrow \nu + \nu$) and neutrino-antineutrino scattering ($\nu + \bar{\nu} \leftrightarrow \nu + \bar{\nu}$) are special cases of the previous electronic neutrino-muonic neutrino scattering but there are a few crucial differences in the derivation of the collision term.

Let us focus first on neutrino autointeractions. It could be viewed as a special case of Eq. (191) in which $a = b = \nu = c = d$. but with two differences. First when examining Eqs. (116) and (117) we realize that this would require to consider $g = G_F/\sqrt{2}$ which is half of the valued used for the interaction between two different species of neutrinos. Since H_I enters as a square, this is a reduction by factor 4. Furthermore, when species a and b are the same species, in the the first line of Eq. (190) we must now use

$$[a_\delta^\dagger a_{\beta'} a_\gamma^\dagger a_{\alpha'}, a_s^\dagger a_{s'}] = A_{\delta\beta'} A_{\gamma s'} \underline{\delta}_{s\alpha'} + A_{\delta s'} A_{\gamma\alpha'} \underline{\delta}_{s\beta'} - A_{\delta\beta'} A_{s\alpha'} \underline{\delta}_{\gamma s'} - A_{s\beta'} A_{\gamma\alpha'} \underline{\delta}_{\delta s'}, \quad (199)$$

that is at this stage of the computation we have twice as many terms and this can be seen as a consequence from the fact that the initial particles are the same [46]. Computing the equivalent of the second line of Eq. (190), using property (30) we find four times more physical terms, which correspond to the product of two allowed pairings in initial states and two in final states. For instance, eliminating the non-physical processes, we find the contribution

$$\langle A_{\beta\delta'} A_{\alpha\gamma'} A_{\delta\beta'} A_{\gamma s'} \rangle \underline{\delta}_{s\alpha'} = (\langle A_{\beta\beta'} \rangle \langle A_{\alpha s'} \rangle + \langle A_{\beta s'} \rangle \langle A_{\alpha\beta'} \rangle) (\langle \widehat{A}_{\alpha\delta'} \rangle \langle \widehat{A}_{\gamma\beta'} \rangle + \langle \widehat{A}_{\alpha\beta'} \rangle \langle \widehat{A}_{\gamma\delta'} \rangle) \underline{\delta}_{s\alpha'}. \quad (200)$$

This factor 4 can also be viewed as the fact that we now have two Feynmann diagrams which add coherently [47]. Taking all factors into account at the level of the collision term [1/4 because the coupling constant g is reduced by half, an extra factor 2 in the first line of Eq. (190) and an extra 4 in the second line of (190)], we end up with an enhancement by a factor 2 compared to neutrino scattering between two different flavors of neutrinos.

Let us focus now on the case of neutrinos scattering on antineutrinos of the same flavor. Again since it comes from the autointeraction of a neutrino flavor neutral current, the coupling constant is also $g = G_F/\sqrt{2}$, that is reduced by a factor 2 compared to scattering between neutrinos and antineutrinos of different flavors. However, when computing

in details the terms which contribute to the Hamiltonian for such process we find 4 terms instead of one. Indeed we recover a double product, so we recover the factor 2 which was initially lost. But we also get twice more terms because there are two Feynmann diagrams for the process (see graphs (a) and (c) of [47]). And with a simple Fierz reordering, which is possible because neutrinos interact with a purely left chiral coupling, they add up coherently. Taking all these factors into account, the interaction Hamiltonian is enhanced by a factor 2 compared to interactions between neutrinos and antineutrinos of different flavors, resulting in a factor 4 enhancement of the collision term.

To summarize, when considering interactions between neutrinos ($\nu + \nu \leftrightarrow \nu + \nu$) one must consider the two-body case (191) in the particular case $a = b = c = d = \nu$ and multiply by a factor 2 the result (this point was omitted in Ref. [48]). And when considering interactions between neutrinos and antineutrinos of the same flavor ($\nu + \bar{\nu} \leftrightarrow \nu + \bar{\nu}$) one must consider the two-body interaction in the particular case $a = c = \nu, b = d = \bar{\nu}$ and multiply the result by a factor 4 in agreement with Ref. [46]. In particular, a simple crossing symmetry allows to get the former reactions $\nu + \nu \leftrightarrow \nu + \nu$ from the $\nu + \bar{\nu} \leftrightarrow \nu + \bar{\nu}$ only up to a factor 1/2. We can interpret this reduction by a factor two using the fact that outgoing particles are identical and one must not double count the outgoing states.

3. neutrino/electron scattering

Contrary to neutrino-neutrino scattering, electron-neutrino scattering is due to both charged and neutral currents. However the Fierz reordering reduces the problem to an interaction of neutral currents with modified chiral couplings. Using Eqs. (120) and (125), the effect of $\nu + e^- \leftrightarrow \nu + e^-$ on neutrinos corresponds to the case

$$a = c = \nu, \quad b = d = e^-, \quad \epsilon_-^{bd} = \epsilon_-^e + 1, \quad \epsilon_+^{bd} = \epsilon_+^e, \quad \epsilon_-^{ac} = e_-^\nu = \frac{1}{2}, \quad \epsilon_+^{ac} = 0, \quad g = 2\frac{G_F}{\sqrt{2}}. \quad (201)$$

The effect of $\nu + e^+ \leftrightarrow \nu + e^+$ is obtained by a crossing symmetry. The effect of $\bar{\nu} + e^+ \leftrightarrow \bar{\nu} + e^+$ on antineutrinos is obtained from charge conjugation of (201), that is it corresponds to the case

$$a = c = \bar{\nu}, \quad b = d = e^+, \quad \epsilon_-^{bd} = \epsilon_-^e + 1, \quad \epsilon_+^{bd} = \epsilon_+^e, \quad \epsilon_-^{ac} = e_-^\nu = \frac{1}{2}, \quad \epsilon_+^{ac} = 0, \quad g = 2\frac{G_F}{\sqrt{2}}, \quad (202)$$

and the effect of $\bar{\nu} + e^- \leftrightarrow \bar{\nu} + e^-$ is obtained from symmetry crossing.

From the electrons perspective, the effect of the $e^- + \nu \leftrightarrow e^- + \nu$ on electrons corresponds to the case

$$b = d = \nu, \quad a = c = e^-, \quad \epsilon_-^{ac} = \epsilon_-^e + 1, \quad \epsilon_+^{ac} = \epsilon_+^e, \quad \epsilon_-^{bd} = e_-^\nu = \frac{1}{2}, \quad \epsilon_+^{bd} = 0, \quad g = 2\frac{G_F}{\sqrt{2}}. \quad (203)$$

and again all other reactions can be obtained from crossing symmetry or charge conjugation as in § VI E.

In all these cases, the generation of linear polarisation has the general form (177). Hence the general conditions to generated polarisation for a given species through two-body interactions are the same. The species under consideration must be massive, and a dipolar structure must be present in either of the two interacting species. For neutrinos, linear polarisation is poorly generated because of their tiny mass and the ratio of the linear polarisation part to the intensity part of the collision term is typically of order m_ν/E_ν . However circular polarisation develops to favor left helicities. For electrons, a linear polarisation can develop, but it is generically erased by the scattering out contributions of the more efficient electromagnetic interactions of the type $e^- + e^- \leftrightarrow e^- + e^-$. Only in some special situations it is necessary to describe correctly the polarisation state of fermions.

Finally, we can check that in the unpolarized case, these results for neutrino/electrons interaction and those for neutrino/neutrinos interactions obtained in § VII B 1 and VII B 2 are exactly the results of Ref. [13]. However note that as mentionned in this reference, there is a typo in the annihilation of neutrino and antineutrinos into electrons and positron in tables 1 and 2 of [46], and thus Tables 1.5 and 1.6 of [49]. The process described in these tables should be of the form $\nu + \bar{\nu} \leftrightarrow e^- + e^+$ and not $\nu + \bar{\nu} \leftrightarrow e^+ + e^-$. Up to this typographical correction our results agree also in the unpolarized case with Refs. [46, 49] and we gather all reactions in Table 3.

C. Majorana neutrinos

In the case where we cannot neglect the mass of neutrinos, there are two possibilities.

- Either the neutrinos are Dirac particles, as we have implicitly assumed so far, and right helicity states are usually underpopulated because the linear polarisation collision term is reduced a factor m_ν/E_ν compared to

Reaction	Particles names	Chiral couplings	$2^{-9} G_F^{-2} E I_{C[A]} = \mathcal{K} \times$
$\nu + \nu_\mu \leftrightarrow \nu + \nu_\mu$	$a + b \leftrightarrow c + d$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \frac{1}{2} \quad \epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})$
$\nu + \bar{\nu}_\mu \leftrightarrow \nu + \bar{\nu}_\mu$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \frac{1}{2} \quad \epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \bar{D})$
$\nu + \bar{\nu} \leftrightarrow \bar{\nu}_\mu + \nu_\mu$	$a + \bar{c} \leftrightarrow \bar{b} + d$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \frac{1}{2} \quad \epsilon_+^{bd} = 0$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, \widehat{C}, D) - \mathcal{T}_I(A, \widehat{B}, \bar{C}, \widehat{D})$
$\nu + \nu \leftrightarrow \nu + \nu$	$a + b \leftrightarrow c + d$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \frac{1}{2} \quad \epsilon_+^{bd} = 0$	$2 [\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})]$
$\nu + \bar{\nu} \leftrightarrow \nu + \bar{\nu}$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \frac{1}{2} \quad \epsilon_+^{bd} = 0$	$4 [\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \bar{D})]$
$\nu + e^- \leftrightarrow \nu + e^-$	$a + b \leftrightarrow c + d$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \epsilon_-^e + 1 \quad \epsilon_+^{bd} = \epsilon_+^e$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, D) - \mathcal{T}_I(A, B, \widehat{C}, \widehat{D})$
$\nu + e^+ \leftrightarrow \nu + e^+$	$a + \bar{d} \leftrightarrow c + \bar{b}$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \epsilon_-^e + 1 \quad \epsilon_+^{bd} = \epsilon_+^e$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, C, \widehat{D}) - \mathcal{T}_I(A, \widehat{B}, \widehat{C}, \bar{D})$
$\nu + \bar{\nu} \leftrightarrow e^+ + e^-$	$a + \bar{c} \leftrightarrow \bar{b} + d$	$\epsilon_-^{ac} = \frac{1}{2} \quad \epsilon_+^{ac} = 0 \quad \epsilon_-^{bd} = \epsilon_-^e + 1 \quad \epsilon_+^{bd} = \epsilon_+^e$	$\mathcal{T}_I(\widehat{A}, \widehat{B}, \widehat{C}, D) - \mathcal{T}_I(A, \widehat{B}, \bar{C}, \widehat{D})$

FIG. 3: Main two-body reactions for the collision term of neutrinos. See Table 2 for more details about the notation. The electronic neutrino is noted ν and the muonic neutrino is noted ν_μ . Similar reactions for antineutrinos can be deduced with a global charge conjugation on all these reactions, and thus on all the collision Kernels.

the intensity part of the collision term. Furthermore to populate the right helicity states (f_{++}) one needs another interaction, according to the structure described at the end of § VIF and this is also reduced by a factor m_ν/E_ν . To a good approximation the quantum neutrino field is excited only in left helicity particles and right helicity antiparticles and the distribution can be considered as being maximally circularly polarized ($I = -V$ for the neutrinos and $I = V$ for the antineutrinos).

- Another possibility is that neutrinos are Majorana particles. In that case they are their own antiparticles. The quantum field (8) is modified by replacing \bar{a}_s^\dagger with a_s^\dagger [20, 50] and consequently in the interaction Hamiltonian we also replace \bar{a}_s by a_s . From Eq. (165a) and using that for a neutral current $J_{aa}^{\mu*} = J_{aa}^\mu$ we see that for a Majorana particle, the effect is that both left and right chiral couplings must be equal in the associated neutral current. That is we must consider that for Majorana neutrinos

$$\epsilon_-^{aa} = \epsilon_+^{aa} = e_-^\nu = \frac{1}{2}. \quad (204)$$

This can be seen more directly by noting from Eqs. (81) and (84) that

$$\bar{v}_r \gamma^\mu (\mathbb{1} - \gamma^5) v_s = \bar{u}_s \gamma^\mu (\mathbb{1} + \gamma^5) u_r. \quad (205)$$

Hence, the collision term for Majorana neutrinos interacting with electrons or positrons are not just the sum of the collision terms of Dirac neutrinos and Dirac antineutrinos. Thanks to CP symmetry, we know already (see § VIE) that the collision term of Dirac antineutrinos can be viewed as a collision term for neutrinos with $\epsilon_+^{ac} = 1/2$ and $\epsilon_-^{ac} = 0$. The interaction of Majorana neutrinos with electrons corresponds to

$$a = c = \nu \quad b = d = e^-, \quad \epsilon_+^{ac} = \epsilon_-^{ac} = e_-^\nu = \frac{1}{2}, \quad \epsilon_-^{bd} = \epsilon_-^e + 1, \quad \epsilon_+^{bd} = \epsilon_+^e, \quad g = 2 \frac{G_F}{\sqrt{2}}. \quad (206)$$

Terms proportional to $(\epsilon_-^{ac})^2$ (resp. $(\epsilon_+^{ac})^2$) are those encountered for Dirac neutrinos (resp. Dirac antineutrinos), and terms proportional to $\epsilon_-^{ac} \epsilon_+^{ac}$ are characteristic of the Majorana nature.

In the case where neutrinos are of Majorana type but massless ($m_a = m_c = 0$), then if they are not linearly polarized, that is if there is no mixing between their Dirac neutrino and Dirac antineutrino nature ($\mathcal{S}_c^{\mu\nu} = \mathcal{S}_a^{\mu\nu} = 0$), then from Eq. (162) we check that this remains the case. Furthermore in the intensity part of the collision term (159), the interference terms proportional to $\epsilon_-^{ac} \epsilon_+^{ac}$ also vanish and their Majorana nature has no physical effect. Finally, still assuming massless Majorana neutrinos interacting with electrons, the circular polarisation part of the collision term (165) remains null if the distribution is initially not polarized, that is if there are as many Dirac neutrinos as Dirac antineutrinos. These conclusions are of course similar when considering interactions with positrons or even autointeractions between neutrinos. To conclude, we recover that in the massless case there is no experiment which can distinguish between Dirac and Majorana neutrinos.

Conclusion

We have shown that the statistical description of fermions is based on a scalar function, the intensity I , and a polarisation vector Q^μ which are both observer independent. The latter encompasses both linear and circular polarisation, but these concepts are observer dependent except in the massless case.

The proper identification of the covariant components I and Q^μ allows to extend the statistical description of fermions to curved space-times, and they should be considered as functions of the position in space-time and of particle momentum. Hence this approach is more general than the more usual description based solely on the distribution functions f_{rs} which is restricted to a Minkowski spacetime, and which requires to consider a given observer to define the helicity states. The connection between these two possible descriptions is made using the spinor valued operator (38) and its decomposition (48) which is a central result of this article.

We then derived a classical Boltzmann equation for the covariant components and we showed in the case of weak interactions that all collision terms are expressed in terms of I and Q^μ . We computed the general collision terms for neutron beta decay and related processes (results gathered in Table 2), and the collision terms for neutrino two-body interactions with other neutrinos or with electrons (results gathered in Table 3), for which we recover the standard results of literature when polarisation is ignored.

The formalism used for the statistical description of fermions is reminiscent of the double projected tensor used to describe photons. However, since linear polarisation for fermions is expanded in spin-1 spherical harmonics, it is generically sourced by distributions with at least a dipolar structure, whereas for photons, linear polarisation is sourced by distribution with at least a quadrupolar dependence as it is expanded in spin-2 spherical harmonics. Furthermore, a second difference is that linear polarisation of fermions is not sourced by the charged current and neutral current couplings if the particle considered is massless.

This difference between fermions and photons can also be understood independently by analyzing the underlying spin structure of the number operators. Since we have combined two spin 1/2 in the number operator of fermions, and thus in their distribution function, it is expected that a spin 1 quantity is involved in the statistical description, and this role is played by the linear polarisation Q^μ part. If fermions are massless, the parallel transport of linear polarisation described by the Liouville equation (92) is in all ways similar to the transport of light polarisation in the eikonal approximation of a light beam. Similarly, when performing the statistical description of photons, we combine two spin-1 quantities in the number operator and thus the distribution function, hence a spin-2 quantity arises as the linear polarisation tensor $\tilde{T}_{\mu\nu}$ [see Appendix. A]. It is no surprise that the description of linear polarisation for a gas of photons is similar to the description of gravitational waves in the eikonal approximation [51]. If in a quantum theory of gravitation, we were to describe statistically the gravitons which are spin-2 particles, we would surely find that their linear polarisation is described by a spin-4 quantity.

In the bosonic case of photons it is known that polarisation is related to the quadrupole moments, but these are typically suppressed in thermal equilibrium. For fermions, polarisation can be generated in thermal equilibrium as it is related to the dipoles and monopoles which cannot be suppressed. We find that both, circular polarisation and linear polarisation appear naturally, but are individually observer dependent.

Finally, we stress that flavor and flavor oscillations can also be described by this formalism, hence it can be used to compute the dynamical evolution of one-particle distribution functions in various contexts, provided the molecular chaos assumption and the classical statistical description hold. However in most cases, linear polarisation is either erased by scattering out processes (e.g. for electrons) or poorly generated (for small mass neutrinos) and the usual approach based only on the intensity I is sufficient.

Acknowledgments

C.F. is supported by the Wallonia-Brussels Federation grant ARC11/15-040 and the Belgian Federal Office for Science, Technical & Cultural Affairs through the Interuniversity Attraction Pole P7/37. C.F. thanks IAP for kind hospitality during the preparation of this article. C.P. is supported by French state funds managed by the ANR within the Investissements d'Avenir programme under reference ANR-11-IDEX-0004-02, and thanks J.-P. Uzan, S. Renaux-Petel and G. Faye for discussions on the kinetic theory.

-
- [1] J. M. Stewart, *Non-Equilibrium Relativistic Kinetic Theory*, vol. 10 (Lecture Notes in Physics, Berlin Springer Verlag, 1971).
 [2] J. Bernstein, *Kinetic theory in the expanding universe* (Cambridge University Press, 1988).

- [3] C.-P. Ma and E. Bertschinger, *Astrophys. J.* **455**, 7 (1995), astro-ph/9506072.
- [4] W. Hu and N. Sugiyama, *Astrophys. J.* **471**, 542 (1996), astro-ph/9510117.
- [5] M. Doran, *JCAP* **0506**, 011 (2005), astro-ph/0503277.
- [6] C. Pitrou, *Phys. Lett.* **B698**, 1 (2011), 1012.0546.
- [7] F.-Y. Cyr-Racine and K. Sigurdson, *Phys. Rev.* **D83**, 103521 (2011), 1012.0569.
- [8] D. Blas, J. Lesgourgues, and T. Tram, *JCAP* **1107**, 034 (2011), 1104.2933.
- [9] D. A. Dicus, E. W. Kolb, A. M. Gleeson, E. C. G. Sudarshan, V. L. Teplitz, and M. S. Turner, *Phys. Rev.* **D26**, 2694 (1982).
- [10] J. Bernstein, L. S. Brown, and G. Feinberg, *Rev. Mod. Phys.* **61**, 25 (1989).
- [11] P. D. Serpico, S. Esposito, F. Iocco, G. Mangano, G. Miele, and O. Pisanti, *JCAP* **0412**, 010 (2004), astro-ph/0408076.
- [12] F. Iocco, G. Mangano, G. Miele, O. Pisanti, and P. D. Serpico, *Phys. Rept.* **472**, 1 (2009), 0809.0631.
- [13] E. Grohs, G. M. Fuller, C. T. Kishimoto, M. W. Paris, and A. Vlasenko, *Phys. Rev.* **D93**, 083522 (2016), 1512.02205.
- [14] K. Kotake, K. Sumiyoshi, S. Yamada, T. Takiwaki, T. Kuroda, Y. Suwa, and H. Nagakura, *PTEP* **2012**, 01A301 (2012), 1205.6284.
- [15] B. Peres, A. J. Penner, J. Novak, and S. Bonazzola, *Class. Quant. Grav.* **31**, 045012 (2014), 1307.1666.
- [16] C. G. Tsagas, A. Challinor, and R. Maartens, *Phys. Rept.* **465**, 61 (2008), 0705.4397.
- [17] C. Pitrou, *Class.Quant.Grav.* **26**, 065006 (2009), 0809.3036.
- [18] D. I. Nagirner and J. Poutanen, *Astron. Astrophys.* **379**, 664 (2001), astro-ph/0108357.
- [19] C. C. Nishi, *Am. J. Phys.* **73**, 1160 (2005), hep-ph/0412245.
- [20] H. K. Dreiner, H. E. Haber, and S. P. Martin, *Phys. Rept.* **494**, 1 (2010), 0812.1594.
- [21] C. Bouchiat and L. Michel, *Muc. Phys.* **5**, 416 (1958).
- [22] A. Vlasenko, G. M. Fuller, and V. Cirigliano, *Phys. Rev.* **D89**, 105004 (2014), 1309.2628.
- [23] D. N. Blaschke and V. Cirigliano (2016), 1605.09383.
- [24] S. Weinberg, *The Quantum Theory of Fields* (Cambridge University Press, 1995).
- [25] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Westview Press, 1995).
- [26] A. Challinor, *Gen. Rel. Grav.* **32**, 1059 (2000), astro-ph/9903283.
- [27] A. Challinor, *Phys. Rev.* **D62**, 043004 (2000), astro-ph/9911481.
- [28] K. S. Thorne, *Rev. Mod. Phys.* **52**, 299 (1980).
- [29] J. N. Goldberg, A. J. Macfarlane, E. T. Newman, F. Rohrlich, and E. C. G. Sudarshan, *J. Math. Phys.* **8**, 2155 (1967).
- [30] R. Durrer, *The Cosmic Microwave Background* (Cambridge University Press, 2008).
- [31] G. Sigl and G. Raffelt, *Nuclear Physics* **B406**, 423 (1993).
- [32] A. Kosowsky, *Annals Phys.* **246**, 49 (1996), astro-ph/9501045.
- [33] M. Beneke and C. Fidler, *Phys. Rev.* **D82**, 063509 (2010), 1003.1834.
- [34] J. Lesgourgues and S. Pastor, *Phys. Rept.* **429**, 307 (2006), astro-ph/0603494.
- [35] J. Lesgourgues and S. Pastor, *Adv. High Energy Phys.* **2012**, 608515 (2012), 1212.6154.
- [36] L. Wolfenstein, *Phys. Rev.* **D17**, 2369 (1978).
- [37] S. P. Mikheev and A. Y. Smirnov, *Nuovo Cim. C* **9**, 17 (1986).
- [38] W. J. Marciano and Z. Parsa, *J. Phys.* **G29**, 2629 (2003), hep-ph/0403168.
- [39] O. Nachtmann and F. Halzen, *Physics Today* **44**, 101 (1991).
- [40] S. Sarantakos, A. Sirlin, and W. J. Marciano, *Nucl. Phys.* **B217**, 84 (1983).
- [41] J. M. Martín-García, *xAct, efficient tensor computer algebra for mathematica* <http://www.xact.es> (2004), URL <http://www.xact.es>.
- [42] R. E. Lopez, M. S. Turner, and G. Gyuk, *Phys. Rev.* **D56**, 3191 (1997), astro-ph/9703065.
- [43] R. E. Lopez and M. S. Turner, *Phys. Rev.* **D59**, 103502 (1999), astro-ph/9807279.
- [44] S. Esposito, G. Mangano, G. Miele, and O. Pisanti, *Nucl. Phys.* **B568**, 421 (2000), astro-ph/9906232.
- [45] F. E. Wietfeldt and G. L. Greene, *Rev. Mod. Phys.* **83**, 1173 (2011).
- [46] A. D. Dolgov, S. H. Hansen, and D. V. Semikoz, *Nucl. Phys.* **B503**, 426 (1997), hep-ph/9703315.
- [47] E. G. Flowers and P. G. Sutherland, *Astrophys. J.* **208**, L19 (1976).
- [48] S. Hannestad and J. Madsen, *Phys. Rev.* **D52**, 1764 (1995), astro-ph/9506015.
- [49] J. Lesgourgues, G. Mangano, G. Miele, and S. Pastor, *Neutrino Cosmology* (Cambridge University Press, 2013).
- [50] P. B. Pal, *Am. J. Phys.* **79**, 485 (2011), 1006.1718.
- [51] L. Blanchet, *Living Reviews in Relativity* **17** (2014), 1310.1528.
- [52] W. Hu and M. J. White, *Phys. Rev.* **D56**, 596 (1997), astro-ph/9702170.
- [53] J. Portsmouth and E. Bertschinger (2004), astro-ph/0412094.

Appendix A: Statistical description for vector bosons

The procedure to build distribution functions is very similar and much simpler for vector bosons. Indeed one starts by considering the number operator (21) and the associated distribution function $f_{rs}(p)$ defined by Eq. (22), but this time with creation and annihilation operators associated with bosons, and which satisfy commutation rules instead of anticommutation rules.

In the Lorentz gauge, for a given momentum p^μ (and given an observer u^μ identified with the time-like vector $[e_0]^\mu$)

of a tetrad), a polarisation basis is given by the three vectors

$$\epsilon_0^\mu(\mathbf{p}) \equiv S^\mu, \quad \epsilon_-^\mu(\mathbf{p}), \quad \epsilon_+^\mu(\mathbf{p}). \quad (\text{A1})$$

which are all unit vectors, mutually orthogonal and transverse to p^μ . A quantum vector field is expanded as

$$\psi(p) = \sum_{s=-1,0,1} \int [d\mathbf{p}] [e^{-ip \cdot x} a_s^\dagger(\mathbf{p}) \epsilon_s^{*\mu}(\mathbf{p}) + e^{ip \cdot x} a_s(\mathbf{p}) \epsilon_s^\mu(\mathbf{p})] \quad (\text{A2})$$

and the quantities which transform like (63), but with the vector representation $D(\Lambda) = \Lambda$, are

$$\sum_{s=-1,0,1} a_s \epsilon_s^\mu, \quad \sum_{s=-1,0,1} a_s^\dagger \epsilon_s^{*\mu}. \quad (\text{A3})$$

A covariant distribution tensor is thus directly obtained by considering

$$f^{\mu\nu}(p) \equiv \sum_{r,s=-1,0,1} f_{rs}(p) \epsilon_r^{*\mu}(\mathbf{p}) \epsilon_s^\nu(\mathbf{p}), \quad (\text{A4})$$

and by construction it is transverse ($f_{\mu\nu} p^\nu = f_{\mu\nu} p^\mu = 0$). Since the spin indices refer to the representation of the rotation group $SU(2)$, for vector bosons the spin indices $-1, 0, 1$ correspond to the irreducible representation $\mathbf{3}$, and the distribution function lives in the (reducible) representation $\mathbf{3} \otimes \mathbf{3}$. By decomposing $f_{\mu\nu}$ in antisymmetric, symmetric traceless, and trace part as

$$f_{\mu\nu}(p) = \frac{1}{2} T_{\mu\nu}(p) - \frac{i}{2m} p^\lambda \epsilon_{\lambda\mu\nu\sigma} V^\sigma(p) + \frac{1}{3} \perp_{\mu\nu} I(p), \quad \perp_{\mu\nu} \equiv \eta_{\mu\nu} + \frac{p_\mu p_\nu}{m^2}, \quad (\text{A5})$$

where $T_{\mu\nu} = T_{(\mu\nu)}$, and with the transverse conditions $T_{\mu\nu} p^\mu = 0$ and $V_\mu p^\mu = 0$, we have decomposed it in irreducible components as $\mathbf{3} \otimes \mathbf{3} = \mathbf{5} \oplus \mathbf{3} \oplus \mathbf{1}$. We recall that the irreducible representations $2\ell + 1$ of the rotation group are spatial symmetric traceless tensor of rank ℓ [28], which here are $T_{\mu\nu}$, V_μ and I .

In order to further decompose the angular dependence, that is the dependence in the direction of the spatial momentum \hat{p}^i , we decompose it according to the sub-representation of the rotation group $SO(2)$ around the helicity direction S^μ according to

$$T_{\mu\nu} = \bar{T} \left(S_\mu S_\nu - \frac{1}{3} \perp_{\mu\nu} \right) + 2S_{(\mu} \bar{T}_{\nu)} + \bar{T}_{\mu\nu}, \quad V_\mu = V S_\mu + \bar{V}_\mu. \quad (\text{A6a})$$

The decomposition in spherical harmonics or spin-weighted spherical harmonics is then

$$I = f_{++} + f_{00} + f_{--}, \quad I = \sum_{\ell m} I_{\ell m}(|\mathbf{p}|) Y_{\ell m}(\hat{\mathbf{p}}) \quad (\text{A7a})$$

$$V = f_{++} - f_{--}, \quad V = \sum_{\ell m} V_{\ell m}(|\mathbf{p}|) Y_{\ell m}(\hat{\mathbf{p}}) \quad (\text{A7b})$$

$$\bar{V}^\mu = \bar{V}^+ \epsilon_+^\mu + \bar{V}^- \epsilon_-^\mu, \quad \bar{V}^+ = f_{-0} - f_{0+}, \quad \bar{V}^- = f_{0-} - f_{+0}, \quad \bar{V}^\pm = \sum_{\ell m} \bar{V}_{\ell m}^\pm(|\mathbf{p}|) Y_{\ell m}^\pm(\hat{\mathbf{p}}) \quad (\text{A7c})$$

$$\bar{T} = 2f_{00} - (f_{++} + f_{--}), \quad \bar{T} = \sum_{\ell m} \bar{T}_{\ell m}(|\mathbf{p}|) Y_{\ell m}(\hat{\mathbf{p}}) \quad (\text{A7d})$$

$$\bar{T}^\mu = \bar{T}^+ \epsilon_+^\mu + \bar{T}^- \epsilon_-^\mu, \quad \bar{T}^\pm = f_{0\pm} + f_{\mp 0}, \quad \bar{T}^\pm = \sum_{\ell m} \bar{T}_{\ell m}^\pm(|\mathbf{p}|) Y_{\ell m}^\pm(\hat{\mathbf{p}}) \quad (\text{A7e})$$

$$\bar{T}^{\mu\nu} = \bar{T}^{++} \epsilon_+^\mu \epsilon_+^\nu + \bar{T}^{--} \epsilon_-^\mu \epsilon_-^\nu, \quad \bar{T}^{\pm\pm} = 2f_{\mp\pm}, \quad \bar{T}^{\pm\pm} = \sum_{\ell m} \bar{T}_{\ell m}^{\pm\pm}(|\mathbf{p}|) Y_{\ell m}^{\pm\pm}(\hat{\mathbf{p}}). \quad (\text{A7f})$$

If the boson vector is massless, as it is the case for photons, then the null-helicity states are not populated because they are not physical so $f_{00} = f_{0\pm} = f_{\pm 0} = 0$, such that $\bar{T} = -I$, $\bar{T}^\pm = \bar{V}^\pm = 0$. For a given observer with four-velocity u^μ , and using the spatial momentum direction vector ($\hat{p}^0 = 0$, $\hat{p}^i = p^i/|\mathbf{p}|$) seen by this observer, the decomposition of the distribution tensor has only four degrees of freedom and reads

$$f_{\mu\nu}(p) = \frac{1}{2} [\bar{T}_{\mu\nu}(p) + \mathcal{H}_{\mu\nu} I(p) - i(u^\lambda \epsilon_{\lambda\mu\nu\sigma} \hat{p}^\sigma) V(p)]. \quad (\text{A8})$$

$I = f_{++} + f_{--}$, $V = f_{++} - f_{--}$ and $\bar{T}^{\pm\pm} \equiv Q \pm iU = 2f_{\mp\pm}$ are the usual Stokes parameters⁵ corresponding to intensity, circular polarisation and linear polarisation. In that case, the distribution tensor is doubly transverse, that is transverse to the momentum p^μ and also to the observer velocity u^μ . In the e_θ, e_ϕ basis its components are the Hermitian 2×2 matrix [16, 30, 52]

$$\frac{1}{2} \begin{pmatrix} I + Q & U - iV \\ U + iV & I - Q \end{pmatrix}. \quad (\text{A9})$$

This structure can also be understood from the fact that when considering the massless representation of the Poincaré group, the little group is no more $SO(3)$ but $ISO(2)$ [24] so the spin structure is classified according to $SO(2) \simeq U(1)$. Hence the number operator and thus the distribution function f_{rs} correspond to the tensor product $\mathbf{2}_1 \otimes \mathbf{2}_1$ (in terms of representations of $SO(2)$) and the latter is decomposed in irreducible representations of $SO(2)$ as $\mathbf{2}_2 \oplus \mathbf{1} \oplus \mathbf{1}$ where $\mathbf{2}_1$ (resp. $\mathbf{2}_2$) is the spin-1 (resp. spin-2) representation of $SO(2)$. In practice and for the physical consequences, the main difference with massless fermions is that linear polarisation is expanded in spin- ± 2 spherical harmonics [52] for massless vector bosons whereas we use spin- ± 1 spherical harmonics for fermions. For massless bosons, linear polarisation is thus sourced by quadrupolar distributions at least (spin- ± 2 spherical harmonics have no monopole nor dipole) whereas for fermions it is sourced by dipolar distributions at least (spin- ± 1 spherical harmonics have no monopole). Finally, just as the collision term for fermions is built out of the covariant parts I and Q^μ , collision terms involving photons are in general built from the double transverse tensor (A8), see e.g. the case of Compton interactions in Refs. [17, 53].

Note also that in all derivations involving bosons, the Pauli blocking operators are everywhere replaced by stimulated emission operators. Adopting the notation $\hat{N}_{rs} \equiv \underline{\delta}_{rs} + N_{rs}$, for the stimulated emission operators, the properties (30) and (B4), which are central in the derivation of the collision term, remain true for bosons. Hence the collision term involving vector bosons, or fermions and bosons, are still formally given by an expression such as (144), see e.g. Eq. (39) of Ref. [33].

Appendix B: Commutation dictionary

We remind the compact notation with multi-indices which for a particle species a is

$$A_{\alpha\alpha'} = N_{s_\alpha s'_\alpha}(p_\alpha, p'_\alpha) = a_{s_\alpha}^\dagger(p_\alpha) a_{s'_\alpha}(p'_\alpha) \quad \underline{\delta}_{\alpha\alpha'} = \delta_{s_\alpha s'_\alpha}^K \underline{\delta}(p_\alpha - p'_\alpha). \quad (\text{B1})$$

From anticommuting rules (9) we get

$$a_{\alpha'} a_\alpha^\dagger = (\underline{\delta}_{\alpha\alpha'} - A_{\alpha\alpha'}) \equiv \hat{A}_{\alpha\alpha'} \quad a_\alpha^\dagger a_{\alpha'} = A_{\alpha\alpha'}. \quad (\text{B2})$$

These basic rules are used to express the results in terms either of the number operator or of the Pauli blocking operator, and since different species commute we can use for instance

$$a_{\alpha'} b_{\beta'} c_{\gamma'} d_\delta^\dagger a_\alpha^\dagger b_\beta^\dagger c_\gamma^\dagger d_{\delta'} = \hat{A}_{\alpha\alpha'} \hat{B}_{\beta\beta'} \hat{C}_{\gamma\gamma'} D_{\delta\delta'}. \quad (\text{B3})$$

It is then immediate to obtain commutators of the form

$$[a_{\alpha'} b_{\beta'}, a_\alpha^\dagger b_\beta^\dagger] = \hat{A}_{\alpha\alpha'} \hat{B}_{\beta\beta'} - A_{\alpha\alpha'} B_{\beta\beta'} \quad (\text{B4a})$$

$$[a_{\alpha'} b_{\beta'} c_{\gamma'} d_{\delta'}, a_\alpha^\dagger b_\beta^\dagger c_\gamma^\dagger d_\delta^\dagger] = \hat{A}_{\alpha\alpha'} \hat{B}_{\beta\beta'} \hat{C}_{\gamma\gamma'} \hat{D}_{\delta\delta'} - A_{\alpha\alpha'} B_{\beta\beta'} C_{\gamma\gamma'} D_{\delta\delta'}. \quad (\text{B4b})$$

For commutators involving strictly more than two operators associated with a given species, we need the commutation rules

$$[a_1, A_{23}] = \underline{\delta}_{12} a_3 \quad [a_1^\dagger, A_{23}] = -\underline{\delta}_{13} a_2^\dagger \quad (\text{B5})$$

$$[A_{12}, A_{34}] = A_{14} \underline{\delta}_{23} - A_{32} \underline{\delta}_{14}, \quad [A_{12} A_{34}, A_{56}] = A_{12} (A_{36} \underline{\delta}_{54} - A_{54} \underline{\delta}_{36}) + A_{34} (A_{16} \underline{\delta}_{52} - A_{52} \underline{\delta}_{16}) \quad (\text{B6})$$

where here 1, 2, 3, 4, 5, 6 represent multi-indices.

⁵ It is sometimes customary in the cosmic microwave background context to define the distribution function as [30] $f_{rs}^{\text{CMB}} \equiv f_{-rs}$. Accordingly, the tensor valued function (A4) is defined as $f^{\mu\nu} = \sum_{rs} f_{rs}^{\text{CMB}} \epsilon_r^\mu \epsilon_s^\nu$. With this definition the Stokes parameters are $I \equiv f_{-++}^{\text{CMB}} + f_{+-}^{\text{CMB}}$, $V \equiv f_{-+-}^{\text{CMB}} - f_{+-}^{\text{CMB}}$ and $Q \pm iU \equiv 2f_{\pm\pm}^{\text{CMB}}$.

Appendix C: Trace technology

The traces of products of γ^μ operators can be found recursively and are non-vanishing only when there is an even number of these operators. Using the identity valid for even n

$$\frac{1}{2}\{\gamma^{\mu_1}, \gamma^{\mu_2} \dots \gamma^{\mu_n}\} = -\sum_{i=2}^n (-1)^i g^{\mu_1 \mu_i} \gamma^{\mu_1} \dots \gamma^{\mu_{i-1}} \gamma^{\mu_{i+1}} \dots \gamma^{\mu_n}, \quad (\text{C1})$$

and the cyclicity of the trace, the first few recursions are

$$T^{\mu\nu} \equiv \text{Tr}[\gamma^\mu \gamma^\nu] = -4g^{\mu\nu} \quad (\text{C2a})$$

$$T^{\mu\nu\rho\sigma} \equiv \text{Tr}[\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma] = -g^{\mu\nu} T^{\rho\sigma} + g^{\mu\rho} T^{\nu\sigma} - g^{\mu\sigma} T^{\nu\rho} = 4(g^{\mu\nu} g^{\rho\sigma} + g^{\mu\sigma} g^{\nu\rho} - g^{\mu\rho} g^{\nu\sigma}) \quad (\text{C2b})$$

$$T^{\mu\nu\rho\sigma\alpha\beta} \equiv \text{Tr}[\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \gamma^\alpha \gamma^\beta] = -g^{\mu\nu} T^{\rho\sigma\alpha\beta} + g^{\mu\rho} T^{\nu\sigma\alpha\beta} - g^{\mu\sigma} T^{\nu\rho\alpha\beta} + g^{\mu\alpha} T^{\nu\rho\sigma\beta} - g^{\mu\beta} T^{\nu\rho\sigma\alpha}. \quad (\text{C2c})$$

When γ^5 matrices are also present in the product, we can simplify it first by gathering them thanks to the anticommutation rule $\gamma^\mu \gamma^5 = -\gamma^5 \gamma^\mu$. If there is an even number of γ^5 , then using $(\gamma^5)^2 = \mathbb{1}$ the problem reduces to a simple product of γ^μ already treated above. If there is an odd number of γ^5 then the problem is reduced to a product of one γ^5 with products of γ^μ . Again the traces of these types of products can be obtained by recursion. Also if there is an odd number of γ^μ or if it is strictly smaller than 4, this type of product vanishes. Using the identity

$$\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho = -\gamma^5 (g^{\mu\nu} \gamma^\rho - g^{\mu\rho} \gamma^\nu + g^{\nu\rho} \gamma^\mu) - i\epsilon^{\mu\nu\rho\sigma} \gamma_\sigma, \quad (\text{C3})$$

the first recursions are

$$T_5^{\mu\nu\rho\sigma} = \text{Tr}[\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma] = 4i\epsilon^{\mu\nu\rho\sigma} \quad (\text{C4a})$$

$$T_5^{\mu\nu\rho\sigma\alpha\beta} = \text{Tr}[\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \gamma^\alpha \gamma^\beta] = -g^{\mu\nu} T_5^{\rho\sigma\alpha\beta} + g^{\mu\rho} T_5^{\nu\sigma\alpha\beta} - g^{\nu\rho} T_5^{\mu\sigma\alpha\beta} - i\epsilon^{\mu\nu\rho\lambda} T^{\lambda\sigma\alpha\beta}. \quad (\text{C4b})$$

The automatic commutation of the γ^5 operators, and the recursive computation of the traces of large products together with their simplifications are very well suited for a computer algebra package and we used *xAct* [41] to deal with these large but otherwise systematic computations.

Appendix D: Two-components spinors

1. Spinors conventions

The two-components spinor formalism is exposed in Ref. [20]. We do not repeat all the definitions in this section and we only gather the useful relations for this article. The two-component spinors indices are raised and lowered with the antisymmetric tensors

$$\epsilon^{\dot{\alpha}\dot{\beta}}, \quad \epsilon^{\alpha\beta}, \quad \epsilon^{\dot{\alpha}\dot{\beta}} = (\epsilon^{\alpha\beta})^* \quad \epsilon^{12} = 1 \quad (\text{D1a})$$

$$\epsilon_{\dot{\alpha}\dot{\beta}}, \quad \epsilon_{\alpha\beta}, \quad \epsilon_{\dot{\alpha}\dot{\beta}} = (\epsilon_{\alpha\beta})^* \quad \epsilon_{12} = -1. \quad (\text{D1b})$$

Indices are conventionally always raised and lowered from the left, for instance as

$$\xi^\alpha = \epsilon^{\alpha\beta} \xi_\beta \quad \xi_\alpha = \epsilon_{\alpha\beta} \xi^\beta, \quad (\text{D2})$$

and the same convention applies for dotted indices. The consequence of this convention is that

$$\epsilon^\alpha{}_\beta = \epsilon^{\alpha\gamma} \epsilon_{\gamma\beta} = \epsilon_{\beta\gamma} \epsilon^{\gamma\alpha} = \epsilon_\beta{}^\alpha = \delta_\beta^\alpha, \quad \epsilon^{\alpha\beta} = -\epsilon^{\alpha\alpha'} \epsilon^{\beta\beta'} \epsilon_{\alpha'\beta'}. \quad (\text{D3})$$

In practice this means that

$$\chi^\alpha \psi_\alpha = -\chi_\alpha \psi^\alpha. \quad (\text{D4})$$

Using the Pauli matrices $\vec{\sigma}$ we define the soldering forms

$$\sigma_{\alpha\dot{\beta}}^\mu = (1, \vec{\sigma}), \quad \sigma_{\mu\alpha\dot{\beta}} = -(1, -\vec{\sigma}) \quad (\text{D5a})$$

$$\bar{\sigma}^{\mu\dot{\alpha}\beta} = (1, -\vec{\sigma}), \quad \bar{\sigma}_\mu^{\dot{\alpha}\beta} = -(1, \vec{\sigma}). \quad (\text{D5b})$$

They are related since

$$\bar{\sigma}^{\mu\dot{\alpha}\beta} = \sigma^{\mu\beta\dot{\alpha}} = \epsilon^{\beta\beta'} \epsilon^{\dot{\alpha}\dot{\alpha}'} \sigma_{\beta'\dot{\alpha}'}^{\mu}. \quad (\text{D6})$$

The two-component spinors are particularly useful because there is a correspondence between vectors and bi-spinors $V_{\alpha\dot{\beta}}$ when using the soldering forms

$$V^{\mu} = \frac{1}{2} \bar{\sigma}^{\mu\dot{\beta}\alpha} V_{\alpha\dot{\beta}}, \quad V_{\alpha\dot{\beta}} = -V^{\mu} \sigma_{\mu\alpha\dot{\beta}}. \quad (\text{D7})$$

These relations can be proved by using the identities

$$\sigma_{\alpha\dot{\alpha}}^{\mu} \bar{\sigma}_{\mu}^{\dot{\beta}\beta} = -2\delta_{\alpha}^{\beta} \delta_{\dot{\alpha}}^{\dot{\beta}} \quad (\text{D8a})$$

$$\sigma_{\alpha\dot{\alpha}}^{\mu} \sigma_{\mu\beta\dot{\beta}} = -2\epsilon_{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \quad (\text{D8b})$$

$$\bar{\sigma}^{\mu\dot{\alpha}\alpha} \bar{\sigma}_{\mu}^{\dot{\beta}\beta} = -2\epsilon^{\alpha\beta} \epsilon^{\dot{\alpha}\dot{\beta}}. \quad (\text{D8c})$$

In particular for a spinor ξ_{α} and its associated $\xi^{\dagger\dot{\alpha}}$, we can define a vector

$$\Xi^{\mu} \equiv \xi^{\alpha} \sigma_{\alpha\dot{\beta}}^{\mu} \xi^{\dagger\dot{\beta}} = \xi_{\dot{\beta}}^{\dagger} \bar{\sigma}^{\mu\dot{\beta}\alpha} \xi_{\alpha} = \xi^{\mu} \xi^{\dagger} = \xi^{\dagger} \bar{\sigma}^{\mu} \xi, \quad (\text{D9})$$

and it has the property to be a null vector $\xi_{\mu} \xi^{\mu} = 0$.

2. Clifford algebra with two-components spinors

A Dirac spinor is expressed in terms of two-components spinors by the decomposition in a direct sum

$$\Psi_{\mathbf{a}} = \begin{pmatrix} \xi_{\alpha} \\ \eta^{\dagger\dot{\alpha}} \end{pmatrix}, \quad (\text{D10})$$

and this relates the Dirac spinor indices \mathbf{a} to the two-component spinor indices. In the chiral representation, the Dirac γ^{μ} matrices are made of 2×2 blocks and read

$$(\gamma^{\mu})_{\mathbf{a}}^{\mathbf{b}} = \begin{pmatrix} 0 & \sigma_{\alpha\dot{\beta}}^{\mu} \\ \bar{\sigma}^{\mu\dot{\alpha}\beta} & 0 \end{pmatrix}. \quad (\text{D11})$$

From the properties

$$[\sigma^{\mu} \bar{\sigma}^{\nu} + \sigma^{\nu} \bar{\sigma}^{\mu}]_{\alpha}^{\beta} = -2g^{\mu\nu} \delta_{\alpha}^{\beta}, \quad [\bar{\sigma}^{\mu} \sigma^{\nu} + \bar{\sigma}^{\nu} \sigma^{\mu}]^{\dot{\alpha}}_{\dot{\beta}} = -2g^{\mu\nu} \delta^{\dot{\alpha}}_{\dot{\beta}} \quad (\text{D12})$$

we check that they satisfy indeed the Clifford algebra

$$\{\gamma^{\mu}, \gamma^{\nu}\} = -2g^{\mu\nu} \begin{pmatrix} \delta_{\alpha}^{\beta} & 0 \\ 0 & \delta^{\dot{\alpha}}_{\dot{\beta}} \end{pmatrix}, \quad (\text{D13})$$

where for simplicity we omit to write the Dirac spinor indices on operators when it is obvious.

To extract chiral components, we also use the matrix

$$\gamma^5 \equiv \frac{i}{4!} \epsilon_{\mu\nu\rho\lambda} \gamma^{\mu} \gamma^{\nu} \gamma^{\rho} \gamma^{\lambda} = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} -\delta_{\alpha}^{\beta} & 0 \\ 0 & \delta^{\dot{\alpha}}_{\dot{\beta}} \end{pmatrix}. \quad (\text{D14})$$

We define also operators in the space of two-components spinors by

$$\sigma^{\mu\nu} = \frac{i}{4} (\sigma^{\mu} \bar{\sigma}^{\nu} - \sigma^{\nu} \bar{\sigma}^{\mu}), \quad \bar{\sigma}^{\mu\nu} = \frac{i}{4} (\bar{\sigma}^{\mu} \sigma^{\nu} - \bar{\sigma}^{\nu} \sigma^{\mu}), \quad (\text{D15})$$

such that the matrices (34) are expressed in blocks as

$$(\Sigma^{\mu\nu})_{\mathbf{a}}^{\mathbf{b}} = \begin{pmatrix} (\sigma^{\mu\nu})_{\alpha}^{\beta} & 0 \\ 0 & (\bar{\sigma}^{\mu\nu})^{\dot{\alpha}}_{\dot{\beta}} \end{pmatrix}. \quad (\text{D16})$$

Since the Dirac matrices are Hermitians, then we find that

$$(\sigma^{\mu\nu})^\dagger = \bar{\sigma}^{\mu\nu}. \quad (\text{D17})$$

Defining the conjugation matrix A and its inverse A^{-1}

$$A^{\text{ab}} = \begin{pmatrix} 0 & \delta_{\dot{\beta}}^{\dot{\alpha}} \\ \delta_{\alpha}^{\beta} & 0 \end{pmatrix} \quad A_{\text{ab}}^{-1} = \begin{pmatrix} 0 & \delta_{\alpha}^{\beta} \\ \delta_{\dot{\beta}}^{\dot{\alpha}} & 0 \end{pmatrix}, \quad (\text{D18})$$

we obtain

$$A \cdot \Sigma^{\mu\nu} \cdot A^{-1} = \Sigma^{\dagger\mu\nu}. \quad (\text{D19})$$

In components and for our chiral representation, we use the numerical equality $A = A^{-1} = \gamma^0$ and it is customary to use the notation γ^0 in place of A . For a Dirac spinor Ψ_{a} decomposed as in Eq. (D10), the Dirac conjugate is defined as

$$\bar{\Psi} \equiv \Psi^\dagger \cdot A \quad \Leftrightarrow \quad \bar{\Psi}^{\text{a}} = \Psi_{\text{b}}^* A^{\text{ba}}. \quad (\text{D20})$$

Finally, the charge conjugation operator defined in § III E 2 is given by

$$C_{\text{ab}} = \begin{pmatrix} \epsilon_{\alpha\beta} & 0 \\ 0 & \epsilon^{\dot{\alpha}\dot{\beta}} \end{pmatrix} \quad (C^{-1})^{\text{ab}} = \begin{pmatrix} \epsilon^{\dot{\alpha}\dot{\beta}} & 0 \\ 0 & \epsilon_{\alpha\beta} \end{pmatrix} \quad (\text{D21})$$

and it satisfies the numerical equality $C = -C^{-1} = i\gamma^0\gamma^2$.

3. Lorentz group representations

An infinitesimal Lorentz transformation is expressed as

$$\Lambda^\mu{}_\nu = \delta^\mu{}_\nu + \omega^\mu{}_\nu, \quad \omega_{\mu\nu} = -\omega_{\nu\mu}. \quad (\text{D22})$$

A finite transformation is obtained through exponentiation of the representation. For Dirac spinors, the representation is obtained as

$$D(\Lambda) = \exp\left(-\frac{i}{2}\omega_{\mu\nu}\Sigma^{\mu\nu}\right), \quad (\text{D23})$$

where we remind the definition (34). In explicit terms, using the two-component spinor expression (D16), it is given by

$$D(\Lambda) = \begin{pmatrix} M & 0 \\ 0 & (M^\dagger)^{-1} \end{pmatrix} \quad M = \exp(-\frac{i}{2}\omega_{\mu\nu}\sigma^{\mu\nu}) \quad (M^\dagger)^{-1} = \exp(-\frac{i}{2}\omega_{\mu\nu}\bar{\sigma}^{\mu\nu}). \quad (\text{D24})$$

The inverse representation is related thanks to the property (D19) which gives

$$D(\Lambda^{-1}) = A^{-1} \cdot D^\dagger(\Lambda) \cdot A, \quad \Leftrightarrow \quad D^{\dagger\text{a}}{}_{\text{c}}(\Lambda) A^{\text{cb}} = A^{\text{ac}} D_{\text{c}}{}^{\text{b}}(\Lambda^{-1}). \quad (\text{D25})$$

The representation M is an element of $\text{SL}(2)$, the covering group of the Lorentz group, and thus conservation of unit determinant implies the property

$$M_{\alpha}{}^{\beta} M_{\alpha'}{}^{\beta'} \epsilon_{\beta\beta'} = \epsilon_{\alpha\alpha'} \quad \Rightarrow \quad M^{-1}{}_{\alpha}{}^{\beta} = -M^{\beta}{}_{\alpha} = -M^T{}_{\alpha}{}^{\beta}. \quad (\text{D26})$$

A left handed two-components spinor ξ_{α} transforms as $M_{\alpha}{}^{\beta}\xi_{\beta}$ under the active transformation Λ . Using (D26), a right handed two-component spinor $\xi^{\dagger\dot{\alpha}}$ transforms as $-M^{\dot{\alpha}}{}_{\dot{\beta}}\xi^{\dagger\dot{\beta}} = (M^{\dagger-1})^{\dot{\alpha}}{}_{\dot{\beta}}\xi^{\dagger\dot{\beta}}$. The representation (D24) acts on a space which is the direct sum of left-handed and right handed two-components spinors in agreement with the decomposition (D10).

When finding explicit expressions for the representation (that is for M and $(M^\dagger)^{-1}$) we have to introduce the notation

$$[\sqrt{-p \cdot \sigma}]_\alpha^\beta \equiv [\sqrt{-p \cdot \sigma \bar{\sigma}^0}]_\alpha^\beta = \frac{-p \cdot \sigma_{\alpha\dot{\gamma}} \bar{\sigma}^{0\dot{\gamma}\beta} + m \delta_\alpha^\beta}{\sqrt{2(E+m)}} \quad (\text{D27a})$$

$$[\sqrt{-p \cdot \bar{\sigma}}]_{\dot{\beta}}^{\dot{\alpha}} \equiv [\sqrt{-p \cdot \bar{\sigma} \sigma^0}]_{\dot{\beta}}^{\dot{\alpha}} = \frac{-p \cdot \bar{\sigma}^{\dot{\alpha}\gamma} \sigma_{\gamma\dot{\beta}}^0 + m \delta_{\dot{\beta}}^{\dot{\alpha}}}{\sqrt{2(E+m)}} \quad (\text{D27b})$$

$$[\sqrt{-p \cdot \sigma}]_{\dot{\beta}}^{\dot{\alpha}} \equiv [\sqrt{-\sigma^0 p \cdot \sigma}]_{\dot{\beta}}^{\dot{\alpha}} = \frac{-\bar{\sigma}^{0\dot{\alpha}\gamma} p \cdot \sigma_{\gamma\dot{\beta}} + m \delta_{\dot{\beta}}^{\dot{\alpha}}}{\sqrt{2(E+m)}} \quad (\text{D27c})$$

$$[\sqrt{-p \cdot \bar{\sigma}}]_\alpha^\beta \equiv [\sqrt{-\sigma^0 p \cdot \bar{\sigma}}]_\alpha^\beta = \frac{-\sigma_{\alpha\dot{\gamma}}^0 p \cdot \bar{\sigma}^{\dot{\gamma}\beta} + m \delta_\alpha^\beta}{\sqrt{2(E+m)}}. \quad (\text{D27d})$$

For detailed expressions see Eqs. (2.106) and (2.107) of Ref. [20]. They satisfy the properties

$$[\sqrt{-p \cdot \sigma}]_\beta^\alpha = -[\sqrt{-p \cdot \bar{\sigma}}]_\beta^\alpha \quad [\sqrt{-p \cdot \bar{\sigma}}]_{\dot{\alpha}}^{\dot{\beta}} = -[\sqrt{-p \cdot \sigma}]_{\dot{\alpha}}^{\dot{\beta}}, \quad (\text{D28})$$

$$\bar{\sigma}^{0\dot{\alpha}\alpha} [\sqrt{-p \cdot \sigma}]_\alpha^\beta \sigma_{\beta\dot{\beta}}^0 = [\sqrt{-p \cdot \sigma}]_{\dot{\beta}}^{\dot{\alpha}}, \quad \bar{\sigma}^{0\dot{\alpha}\alpha} [\sqrt{-p \cdot \bar{\sigma}}]_\alpha^\beta \sigma_{\beta\dot{\beta}}^0 = [\sqrt{-p \cdot \bar{\sigma}}]_{\dot{\beta}}^{\dot{\alpha}}, \quad (\text{D29})$$

which can be used to change the position of indices. The definitions (D27) are convenient since it can then be shown that for a pure boost which transforms a momentum from its rest-frame to a frame where it takes on-shell values p^μ , the representations are

$$M = \sqrt{\frac{-p \cdot \sigma}{m}}, \quad (M^\dagger)^{-1} = \sqrt{\frac{-p \cdot \bar{\sigma}}{m}}. \quad (\text{D30})$$

4. Plane wave solutions of the Dirac equation

In two-components form, the wave-functions u_s and v_s are of the form

$$u_{s,\mathbf{a}}(\mathbf{p}) = \begin{pmatrix} x_{s,\alpha}(\mathbf{p}) \\ y_s^{\dagger\dot{\alpha}}(\mathbf{p}) \end{pmatrix} \quad v_{s,\mathbf{a}}(\mathbf{p}) = \begin{pmatrix} y_{s,\alpha}(\mathbf{p}) \\ x_s^{\dagger\dot{\alpha}}(\mathbf{p}) \end{pmatrix}. \quad (\text{D31})$$

To obtain explicit expressions, let us take a basis for the two-components spinors. This basis will be used for both the two-components spinors with down index and dotted up index. For down indices we note this basis $(\chi_s)_\alpha$ and for dotted up indices $(\eta_s^\dagger)^{\dot{\alpha}}$. Using a direction $\hat{\mathbf{s}}$ having angles θ, ϕ in spherical coordinates, the components of this basis are

$$\left(\chi_{\frac{1}{2}}\right)_\alpha = \left(\eta_{\frac{1}{2}}^\dagger\right)^{\dot{\alpha}} = \begin{pmatrix} e^{-i\phi/2} \cos(\theta/2) \\ e^{i\phi/2} \sin(\theta/2) \end{pmatrix} = e^{-\frac{i}{2}\phi\sigma_z} \cdot e^{-\frac{i}{2}\theta\sigma_y} \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{D32a})$$

$$\left(\chi_{-\frac{1}{2}}\right)_\alpha = \left(\eta_{-\frac{1}{2}}^\dagger\right)^{\dot{\alpha}} = \begin{pmatrix} -e^{-i\phi/2} \sin(\theta/2) \\ e^{i\phi/2} \cos(\theta/2) \end{pmatrix} = e^{-\frac{i}{2}\phi\sigma_z} \cdot e^{-\frac{i}{2}\theta\sigma_y} \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (\text{D32b})$$

The building blocks of u_s and v_s are then given by

$$x_{s,\alpha}(\mathbf{p}) = [\sqrt{p \cdot \sigma}]_\alpha^\beta (\chi_s)_\beta \quad y_s^{\dagger\dot{\alpha}}(\mathbf{p}) = [\sqrt{p \cdot \bar{\sigma}}]_{\dot{\beta}}^{\dot{\alpha}} (\eta_s^\dagger)^{\dot{\beta}}, \quad (\text{D33a})$$

$$x_s^{\dagger\dot{\alpha}}(\mathbf{p}) = (\chi_s^\dagger)_{\dot{\beta}} [\sqrt{p \cdot \sigma}]_{\dot{\beta}}^{\dot{\alpha}} \quad y_{s,\alpha}(\mathbf{p}) = (\eta_s)^\beta [\sqrt{p \cdot \bar{\sigma}}]_\beta^\alpha. \quad (\text{D33b})$$

Raising and lowering indices on these previous expressions and using properties (D28) we find

$$x_s^\alpha(\mathbf{p}) = (\chi_s)^\beta [\sqrt{p \cdot \bar{\sigma}}]_\beta^\alpha \quad y_{s,\dot{\alpha}}^\dagger(\mathbf{p}) = (\eta_s^\dagger)_{\dot{\beta}} [\sqrt{p \cdot \sigma}]_{\dot{\beta}}^{\dot{\alpha}} \quad (\text{D34a})$$

$$x_s^{\dagger\dot{\alpha}}(\mathbf{p}) = [\sqrt{p \cdot \bar{\sigma}}]_{\dot{\beta}}^{\dot{\alpha}} (\chi_s^\dagger)^{\dot{\beta}} \quad y_{s,\alpha}(\mathbf{p}) = [\sqrt{p \cdot \sigma}]_\alpha^\beta (\eta_s)_\beta. \quad (\text{D34b})$$

In order to compare with Eqs. (3.1.19-3.1.22) of Ref. [20] we must use the relations

$$(\chi_s)^\alpha = -2s(\eta_{-s})^\alpha \quad (\chi_s^\dagger)^{\dot{\alpha}} = -2s(\eta_{-s}^\dagger)^{\dot{\alpha}} \quad (\text{D35a})$$

$$(\eta_s)_\alpha = 2s(\chi_{-s})_\alpha \quad (\eta_s^\dagger)_{\dot{\alpha}} = 2s(\chi_{-s}^\dagger)_{\dot{\alpha}}. \quad (\text{D35b})$$

Eqs. (D33) and (D34) are all what is needed to get explicit expressions for u_s and v_s from Eqs. (D31), but also for $\bar{u}_s = u_s^\dagger \gamma^0$ and $\bar{v}_s = v_s^\dagger \gamma^0$ which are expressed in terms of two-components spinors as

$$\bar{u}_s^\alpha(\mathbf{p}) = \begin{pmatrix} y_s^\alpha(\mathbf{p}) \\ x_{s,\dot{\alpha}}^\dagger(\mathbf{p}) \end{pmatrix} \quad v_s^\alpha(\mathbf{p}) = \begin{pmatrix} x_s^\alpha(\mathbf{p}) \\ y_{s,\dot{\alpha}}^\dagger(\mathbf{p}) \end{pmatrix}. \quad (\text{D36})$$

Note that the plane-wave solutions for a momentum p^μ are obtained simply by considering the case where the momentum is at rest, for which $x_{s,\alpha} = (\chi_s)_\alpha$ and $y_s^{\dagger\dot{\alpha}}(\mathbf{p}) = (\eta_s^\dagger)^{\dot{\alpha}}$ and then boosting using Eqs. (D24) and (D30).

5. Helicity basis plane wave solutions

If the direction $\hat{\mathbf{s}}$ used to build the basis is the spatial momentum direction $\hat{\mathbf{p}} = (\sin(\theta)\cos(\phi), \sin(\theta)\sin(\phi), \cos(\theta))$, then we are dealing with helicities and we have

$$\frac{1}{2}\hat{\mathbf{s}} \cdot \vec{\sigma}\chi_s = s\chi_s \quad \frac{1}{2}\hat{\mathbf{s}} \cdot \vec{\sigma}\eta_s = s\eta_s. \quad (\text{D37})$$

The relations (D27) take in that case a very simple form, and we obtain the relativistic two-components spinors (D33) immediately

$$x_{s,\alpha}(p) = \sqrt{E - 2s|p|}(\chi_s)_\alpha \quad y_s^{\dagger\dot{\alpha}}(p) = \sqrt{E + 2s|p|}(\eta_s^\dagger)^{\dot{\alpha}} \quad (\text{D38a})$$

$$x_s^{\dagger\dot{\alpha}}(p) = \sqrt{E - 2s|p|}(\chi_s^\dagger)^{\dot{\alpha}} \quad y_{s,\alpha}(p) = \sqrt{E + 2s|p|}(\eta_s)_\alpha. \quad (\text{D38b})$$

Using the general form of the plane wave solutions (D31) and Eqs. (D35) we obtain the helicity solutions (20).

6. Bilinears with two-components spinors

Using the definitions (39) and (42) we obtain bilinear scalars built from two-components spinors

$$X_{++}^\mu \equiv x_{+,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} x_{+,\alpha} = x_+^\alpha \sigma_{\alpha\dot{\beta}}^\mu x_+^{\dagger\dot{\beta}} = (E - |p|)(1, -\hat{p}) \quad (\text{D39a})$$

$$X_{--}^\mu \equiv x_{-,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} x_{-,\alpha} = x_-^\alpha \sigma_{\alpha\dot{\beta}}^\mu x_-^{\dagger\dot{\beta}} = (E + |p|)(1, \hat{p}) \quad (\text{D39b})$$

$$X_{-+}^\mu \equiv x_{-,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} x_{+,\alpha} = x_+^\alpha \sigma_{\alpha\dot{\beta}}^\mu x_-^{\dagger\dot{\beta}} = -\sqrt{2}m(0, \epsilon_-) \quad (\text{D39c})$$

$$X_{+-}^\mu \equiv x_{+,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} x_{-,\alpha} = x_-^\alpha \sigma_{\alpha\dot{\beta}}^\mu x_+^{\dagger\dot{\beta}} = -\sqrt{2}m(0, \epsilon_+) \quad (\text{D39d})$$

$$Y_{++}^\mu \equiv y_{+,\beta}^\alpha \sigma_{\alpha\dot{\beta}}^\mu y_{+,\dot{\beta}}^\dagger = y_{+,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} y_{+,\alpha} = (E + |p|)(1, \hat{p}) \quad (\text{D40a})$$

$$Y_{--}^\mu \equiv y_{-,\beta}^\alpha \sigma_{\alpha\dot{\beta}}^\mu y_{-,\dot{\beta}}^\dagger = y_{-,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} y_{-,\alpha} = (E - |p|)(1, -\hat{p}) \quad (\text{D40b})$$

$$Y_{-+}^\mu \equiv y_{-,\beta}^\alpha \sigma_{\alpha\dot{\beta}}^\mu y_{+,\dot{\beta}}^\dagger = y_{+,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} y_{-,\alpha} = \sqrt{2}m(0, \epsilon_-) \quad (\text{D40c})$$

$$Y_{+-}^\mu \equiv y_{+,\beta}^\alpha \sigma_{\alpha\dot{\beta}}^\mu y_{-,\dot{\beta}}^\dagger = y_{-,\beta}^\dagger \bar{\sigma}^{\mu\dot{\beta}\alpha} y_{+,\alpha} = \sqrt{2}m(0, \epsilon_+) \quad (\text{D40d})$$

$$(YX)_{++}^{0i} \equiv y_{+,\alpha}^\dagger [\sigma^{0i}]_\alpha^\beta x_{+,\beta} = x_{+,\alpha}^\dagger [\sigma^{0i}]_\alpha^\beta y_{+,\beta} = -\frac{im}{2}\hat{p}^i \quad (\text{D41a})$$

$$(YX)_{++}^{ij} \equiv y_{+,\alpha}^\dagger [\sigma^{ij}]_\alpha^\beta x_{+,\beta} = x_{+,\alpha}^\dagger [\sigma^{ij}]_\alpha^\beta y_{+,\beta} = \epsilon^{ij}_k \frac{m}{2}\hat{p}^k \quad (\text{D41b})$$

$$(YX)_{--}^{0i} \equiv y_{-,\alpha}^\dagger [\sigma^{0i}]_\alpha^\beta x_{-,\beta} = x_{-,\alpha}^\dagger [\sigma^{0i}]_\alpha^\beta y_{-,\beta} = \frac{im}{2}\hat{p}^i \quad (\text{D41c})$$

$$(YX)_{--}^{ij} \equiv y_{-,\alpha}^\dagger [\sigma^{ij}]_\alpha^\beta x_{-,\beta} = x_{-,\alpha}^\dagger [\sigma^{ij}]_\alpha^\beta y_{-,\beta} = -\epsilon^{ij}_k \frac{m}{2}\hat{p}^k \quad (\text{D41d})$$

$$(YX)_{-+}^{0i} \equiv y_-^\alpha [\sigma^{0i}]_\alpha^\beta x_{+,\beta} = x_+^\alpha [\sigma^{0i}]_\alpha^\beta y_{-,\beta} = -\frac{i}{\sqrt{2}}(E - |p|)\epsilon_-^i \quad (\text{D42a})$$

$$(YX)_{-+}^{ij} \equiv y_-^\alpha [\sigma^{ij}]_\alpha^\beta x_{+,\beta} = x_+^\alpha [\sigma^{ij}]_\alpha^\beta y_{-,\beta} = \frac{1}{\sqrt{2}}(E - |p|)\epsilon_{-k}^i \epsilon_{-l}^j \quad (\text{D42b})$$

$$(YX)_{+-}^{0i} \equiv y_+^\alpha [\sigma^{0i}]_\alpha^\beta x_{-,\beta} = x_-^\alpha [\sigma^{0i}]_\alpha^\beta y_{+,\beta} = -\frac{i}{\sqrt{2}}(E + |p|)\epsilon_+^i \quad (\text{D42c})$$

$$(YX)_{+-}^{ij} \equiv y_+^\alpha [\sigma^{ij}]_\alpha^\beta x_{-,\beta} = x_-^\alpha [\sigma^{ij}]_\alpha^\beta y_{+,\beta} = \frac{1}{\sqrt{2}}(E + |p|)\epsilon_{+k}^i \epsilon_{+l}^j \quad (\text{D42d})$$

$$(XY)_{++}^{0i} \equiv x_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha y_{+,\dot{\beta}}^\dagger = y_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha x_{+,\dot{\beta}}^\dagger = \frac{im}{2}\hat{p}^i \quad (\text{D43a})$$

$$(XY)_{++}^{ij} \equiv x_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha y_{+,\dot{\beta}}^\dagger = y_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha x_{+,\dot{\beta}}^\dagger = \epsilon_{-k}^{ij} \frac{m}{2}\hat{p}^k \quad (\text{D43b})$$

$$(XY)_{--}^{0i} \equiv x_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha y_{-,\dot{\beta}}^\dagger = y_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha x_{-,\dot{\beta}}^\dagger = -\frac{im}{2}\hat{p}^i \quad (\text{D43c})$$

$$(XY)_{--}^{ij} \equiv x_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha y_{-,\dot{\beta}}^\dagger = y_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha x_{-,\dot{\beta}}^\dagger = -\epsilon_{-k}^{ij} \frac{m}{2}\hat{p}^k \quad (\text{D43d})$$

$$(XY)_{-+}^{0i} \equiv x_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha y_{+,\dot{\beta}}^\dagger = y_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha x_{-,\dot{\beta}}^\dagger = \frac{i}{\sqrt{2}}(E + |p|)\epsilon_-^i \quad (\text{D44a})$$

$$(XY)_{-+}^{ij} \equiv x_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha y_{+,\dot{\beta}}^\dagger = y_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha x_{-,\dot{\beta}}^\dagger = \frac{1}{\sqrt{2}}(E + |p|)\epsilon_{-k}^i \epsilon_{-l}^j \quad (\text{D44b})$$

$$(XY)_{+-}^{0i} \equiv x_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha y_{-,\dot{\beta}}^\dagger = y_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{0i}]_{\dot{\beta}}^\alpha x_{+,\dot{\beta}}^\dagger = \frac{i}{\sqrt{2}}(E - |p|)\epsilon_+^i \quad (\text{D44c})$$

$$(XY)_{+-}^{ij} \equiv x_{+,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha y_{-,\dot{\beta}}^\dagger = y_{-,\dot{\alpha}}^\dagger [\bar{\sigma}^{ij}]_{\dot{\beta}}^\alpha x_{+,\dot{\beta}}^\dagger = \frac{1}{\sqrt{2}}(E - |p|)\epsilon_{+k}^i \epsilon_{+l}^j \quad (\text{D44d})$$

7. Bilinears with Dirac spinors

From the explicit expression of the plane wave solutions in terms of two-components spinors (D31) and (D36), and the explicit expression of the $\gamma^\mu, \gamma^5, \Sigma^{\mu\nu}$ matrices of § D 2, we find the needed scalar bilinear combinations

$$\bar{u}_r u_s = 2m\delta_{rs}^K \quad (\text{D45a})$$

$$\bar{u}_r \gamma^5 u_s = 0 \quad (\text{D45b})$$

$$\bar{u}_r \gamma^\mu u_s = X_{rs}^\mu + Y_{rs}^\mu = 2\delta_{rs}^K p^\mu \quad (\text{D45c})$$

$$\bar{u}_r \gamma^\mu \gamma^5 u_s = -X_{rs}^\mu + Y_{rs}^\mu = \delta_{rs}^K (4rmS^\mu) + \delta_{r-s}^K \left(-\sqrt{8}m\epsilon_{s-r}^\mu \right) \quad (\text{D45d})$$

$$\bar{u}_r \tilde{\Sigma}^{0i} u_s = -i(YX)_{rs}^{0i} + i(XY)_{rs}^{0i} = \delta_{rs}^K (-2rm\hat{p}^i) + \delta_{r-s}^K \left(-\sqrt{2}E\epsilon_{r-s}^i \right) \quad (\text{D45e})$$

$$\bar{u}_r \tilde{\Sigma}^{ij} u_s = -i(YX)_{rs}^{ij} + i(XY)_{rs}^{ij} = \delta_{r-s}^K \left(\sqrt{2}i(s-r)|\hat{p}| \epsilon_{r-s}^i \epsilon_{r-s}^j \right) = \delta_{r-s}^K \left(-2\sqrt{2}p^{[i} \epsilon_{r-s}^{j]} \right), \quad (\text{D45f})$$

where δ_{r-s}^K vanishes if $r = s$ and is unity otherwise. When forming the similar bilinears but with negative frequency plane-wave solutions $v_s(p)$ we find

$$\bar{v}_r v_s = -2m\delta_{rs}^K \quad (\text{D46a})$$

$$\bar{v}_r \gamma^5 v_s = 0 \quad (\text{D46b})$$

$$\bar{v}_r \gamma^\mu v_s = Y_{sr}^\mu + X_{sr}^\mu = 2\delta_{rs}^K p^\mu \quad (\text{D46c})$$

$$\bar{v}_r \gamma^\mu \gamma^5 v_s = -Y_{sr}^\mu + X_{sr}^\mu = \delta_{rs}^K (-4rmS^\mu) + \delta_{r-s}^K \left(\sqrt{8}m\epsilon_{s-r}^\mu \right) \quad (\text{D46d})$$

$$\bar{v}_r \tilde{\Sigma}^{0i} v_s = -i(XY)_{sr}^{0i} + i(YX)_{sr}^{0i} = \delta_{rs}^K (2rm\hat{p}^i) + \delta_{r-s}^K \left(-\sqrt{2}E\epsilon_{r-s}^i \right) \quad (\text{D46e})$$

$$\bar{v}_r \tilde{\Sigma}^{ij} v_s = -i(XY)_{sr}^{ij} + i(YX)_{sr}^{ij} = \delta_{r-s}^K \left(-2\sqrt{2}p^{[i} \epsilon_{r-s}^{j]} \right). \quad (\text{D46f})$$

We realize that the bilinears built from \bar{v}_r and v_s are obtained from those built with \bar{u}_r and u_s by the simple replacement $m \rightarrow -m$.

8. Covariant components of the distribution function using two-components spinors

The covariant components of the distribution function f_{rs} can also be extracted with the two-components formalism, and it happens to lead to straightforward expressions. Let us define first the distribution functions which are 2×2 valued in complex spinor space by

$$f_{\dot{\alpha}\beta}^x(p) \equiv \sum_{rs} x_{r,\dot{\alpha}}^\dagger(p) f_{rs}(p) x_{s,\beta}(p) \quad f_y^{\alpha\dot{\beta}}(p) \equiv \sum_{rs} y_r^\alpha(p) f_{rs}(p) y_s^{\dot{\beta}\dagger}(p). \quad (\text{D47})$$

Due to the relations (D7), we can define 4-vector distribution functions simply as

$$f_x^\mu(p) = \frac{1}{2} \bar{\sigma}^{\mu\dot{\alpha}\beta} f_{\dot{\alpha}\beta}^x(p), \quad f_y^\mu(p) = \frac{1}{2} \sigma_{\alpha\dot{\beta}}^\mu f_y^{\alpha\dot{\beta}}(p). \quad (\text{D48})$$

Using the definitions (46) and (47) we get

$$f_x^\mu = \frac{1}{2} (Ip^\mu - mQ^\mu) = \begin{cases} \mathcal{I}^{-\mu} & \text{particles} \\ \mathcal{I}^{+\mu} & \text{antiparticles,} \end{cases} \quad f_y^\mu = \frac{1}{2} (Ip^\mu + mQ^\mu) = \begin{cases} \mathcal{I}^{+\mu} & \text{particles} \\ \mathcal{I}^{-\mu} & \text{antiparticles.} \end{cases} \quad (\text{D49})$$

The interpretation is thus that \mathcal{I}_μ^\pm are vectors of chiral parts, with \mathcal{I}_μ^- (resp. \mathcal{I}_μ^+) being a the left chiral part (resp. right chiral part) vector for particles or the right chiral part (resp. left chiral part) vector for antiparticles.